



Sanford Hillsboro
2016 Community Health
Needs Assessment

SANFORD[®]
HEALTH

Sanford Hillsboro Medical Center

Community Health Needs Assessment

2016

Dear Community Members,

Sanford Hillsboro is pleased to present the 2016 Community Health Needs Assessment.

Part of the comprehensive assessment work is to formally identify unmet health needs in the community. Community stakeholders helped to prioritize the unmet needs. The prioritized needs will be a focus for strategic planning and the required implementation strategy development. We are grateful to all the public health and community members who joined us in this important work.

During 2015 members of the community were asked to complete a survey to help identify unmet health needs. Researchers at the Center for Social Research at North Dakota State University analyzed the survey data. Sanford further analyzed the data, identified unmet needs, and partnered with key community stakeholders to develop a list of resources and assets that were available to address each need. A gap analysis and prioritization exercise was also conducted to identify the most significant health needs and to further address these needs through the implementation strategies that are included in this document.

Sanford Hillsboro will focus on strategies to address and improve the following community health needs:

- Physical Health
- Mental Health Services

The report focuses on community assets as well as community health needs. The asset map/resource list is included in this document along with the action steps that will be taken to address each identified need.

At Sanford Hillsboro, patient care extends beyond our bricks and mortar. As a not-for-profit organization, ensuring that the benefits of health care reach the broad needs of communities is at the core of who we are. Through our work with communities, we can bring health and healing to the people who live and work across our communities. Together, we can fulfill this mission.

Sincerely,



Jac McTaggart
Chief Executive Officer
Sanford Medical Center Hillsboro

Sanford Hillsboro Medical Center
Community Health Needs Assessment
2016

EXECUTIVE SUMMARY

Sanford Hillsboro Medical Center

Community Health Needs Assessment 2016

Purpose

A community health needs assessment is critical to a vital community benefit program that builds on community assets, promotes collaboration, improves community health, and promotes innovation and research. A community health needs assessment helps the community build capacity to support policy, systems, environmental changes and community health improvement. A community health needs assessment also serves to validate progress made toward organizational strategies and provides further evidence for retaining not-for-profit status.

The purpose of this community health needs assessment is to develop a global view of the population's health and the prevalence of disease and health issues within our community. Findings from the assessment serve as a catalyst to align expertise and to develop a community benefit plan of action. There is great intrinsic value in a community health needs assessment when it serves to validate, justify and defend not-for-profit status and create opportunity to identify and address public health issues from a broad perspective.

Study Design and Methodology

1. Non-Generalizable Survey

An on-line non-generalizable survey was conducted on-line through a partnership between Sanford Hillsboro, Sanford Mayville, Traill County Public Health, Steele County Public Health and the City of Halstad and the Center for Social Research (CSR) at North Dakota State University. The CSR developed and maintained links to the on-line survey tool. The website address for the survey instrument was distributed via e-mail to various agencies, at times using a snowball approach. Data collection occurred throughout the month of April 2015 and a total of 54 respondents participated in the on-line survey.

The purpose of this non-generalizable survey of community leaders was to learn about the perceptions of area community leaders regarding community health, their personal health, preventive health, and the prevalence of disease. This group included community leaders, legislators, and agency leaders representing chronic disease and disparity.

A Likert scale was developed to determine the respondent's highest concerns, with 1 as not at all and 5 meaning a great deal. Needs ranking 3.5 and above were included in the needs to be addressed and prioritized. Many of the identified needs that ranked below 3.5 are being addressed by Sanford and community partners. However, 3.5 and above was used as a focus for the purpose of the required prioritization.

2. Community Stakeholder Meeting

Community stakeholders were invited to a meeting to review the early findings from the survey and to discuss the top health issues or health-related issues facing the community. The attendees at the September 2015 community stakeholder meeting helped to determine key priorities for the community.

3. Community Asset Mapping

Asset mapping was conducted by reviewing the data and identifying the unmet needs from the various surveys and data sets. Each unmet need was researched to determine what resources were available in the community to address the needs. Once gaps were determined the group proceeded to the prioritization process. A multi-voting methodology was implemented to determine what top priorities would be further developed into implementation strategies.

4. Secondary Research

The secondary data includes community health profiles from the North Dakota Department (DOH) of Health for Traill and Steele Counties, North Dakota and the Robert Wood Johnson County Health Rankings. The (DOH) indicators that were reviewed for this assessment include: population data, vital statistics, adult behavioral risk factors, crime and child risk.

Key Findings – Primary Research

The key findings are based on the generalizable and the non-generalizable survey data. Key indicators were ranked on a 1-5 Likert scale, with 5 being the highest concern ranking. Survey results that rank 3.5 or higher are considered to be high ranking concerns the survey.

1. **Aging:** The number one ranking concern among respondents overall is the cost of long term care. The availability of memory care and the availability of long term care also rank as top concerns for the aging.
2. **Children and Youth:** When considering children and youth, the availability and cost of activities for youth, availability and cost of quality childcare, bullying, the availability and cost of quality infant care, and the availability and cost of services for at-risk youth, are also ranked as high concerns.
3. **Safety:** The presence of street drugs and alcohol and drug dealers in the community, child abuse and neglect, and domestic violence are the highest safety concerns of the respondents.

4. **Health Care:** The health care indicator addressed access to affordable health insurance, access to affordable health care, the cost of affordable dental insurance, and the cost of affordable vision insurance, timely access to substance abuse providers and mental health providers, and affordable prescription drug coverage are the highest concerns among the respondents.
5. **Physical Health:** Cancer, chronic disease, inactivity and lack of exercise, obesity, and poor nutrition and eating habits are the highest physical health concerns.
6. **Mental Health/Behavioral Health:** Depression, stress, underage drug use and abuse, underage drinking, drug use and abuse, smoking and tobacco, and alcohol use and abuse are the highest concerns for mental health/behavioral health.

Key Findings – Secondary Research Based on the 2015 County Health Rankings

Health Outcomes

Premature death: The premature death indicator is defined as years of potential life lost before age 75 per 100,000 population. The mortality health outcomes indicate that North Dakota as a state has more premature deaths (307.1 per 100,000) than Minnesota (262.6 per 100,000).

Trail County, North Dakota was ranked best in the state as #1 out of 47 counties for length of life. The premature deaths in Trail County (4,400) rate much lower than the state of North Dakota at 6,388.

Poor or fair health: 13% of adults in Trail County and 12% in Steele County report poor or fair health compared to 10% for the nation, 12% in North Dakota, and 11% in Minnesota.

The average number of days reported in the last 30 as unhealthy mental health days is 2.2 in Trail County and 2.3 in Steele County. The state of North Dakota reports 2.4 days, and Minnesota reports 2.6 days.

The percent of live births with low birth weight (less than 2,500 grams) is 4.2% in Trail County and there is not enough data to rank for Steele County. Both North Dakota and Minnesota report 6.5%.

Health Factors

The percent of adults who are currently smoking is 13% in Steele County and 12% in Steele County. The state of North Dakota reports 18% of adults are current smokers and in Minnesota the percent of current adult smokers is 16%.

36% of the adult population in Trail County, and 32% in Steele County are considered obese with a BMI over 30. This compares with North Dakota at 30% and Minnesota at 26%.

The percent of adults reporting excessive or binge drinking is 16% in Trail County and 26% in Steele County. North Dakota reports 22% statewide and Minnesota reports 19%.

Driving deaths that have alcohol involvement is at 33% in Trail County and 50% in Steele County. North Dakota as a whole is at 46% and Minnesota is at 31%.

Sexually transmitted infections rank substantially higher than the national benchmark (138) for North Dakota (416), Minnesota (336), and Traill (161); however, Steele County is not ranked. The teen birth rate is higher in North Dakota (28) and Minnesota (24) than the national benchmark (20), but is lower in Traill (15) and slightly higher in Steele County (21).

The clinical care outcomes indicate that North Dakota and Steele County (12%) have a higher percentage of uninsured adults, and Traill County is at the same rate and the national benchmark.

The ratio of population to primary care physicians is 1,279:1 in North Dakota and 1,113:1 in Minnesota. Traill County's ratio is 4,036:1, while Steele County's ratio is more favorable at 1,989:1.

The ratio of population to mental health providers is 638:1 in North Dakota and 529:1 in Minnesota. Traill and Steele Counties are not ranked. The number of professionally active dentists is 1,710:1 North Dakota, 4,123:1 in Traill County, and Steele County is not ranked.

Preventable hospital stays in Traill County are at 55; and in North Dakota 56. Diabetic screening is at 92% in Traill County and 86% in North Dakota. Steele County is slightly lower at 87%. Mammography screening is at 74.3% in Traill County and 78.9% in Steele County compared to 68% in North Dakota.

The social and economic factor outcomes indicate that North Dakota has a high school graduate rate of 85%. The high school graduation rate is not available for Traill and Steele counties. Post-secondary education (some post-secondary education) is at 76.4% in Traill County, 72.3% in Steele County and 74.4% in North Dakota.

The unemployment rate is 2.9% in North Dakota, 3.1% in Traill County, and 2.2% in Steele County. The percentage of child poverty is 12% in North Dakota and 9% in Traill County. Child poverty is 12% in Steele County.

Social associations are defined as the number of membership associations per 10,000 population, and are ranked higher in Traill County at 32.2 than in North Dakota at 17.3. Steele County is lower at 10.1.

The percentage of children in single parent households is 15% in Steele County, 20% in Traill County, and 26% in North Dakota.

Violent crime is lower in Traill County at 48 per 100,000 population than in North Dakota at 240 per 100,000 population. Steele County is at 25 per 100,000 population.

Youth account for 22% of the population in Traill County and 21.5% of the population in Steele County. Elderly account for 18.6% of the population in Traill County and 19% of the population in Steele County.

The following needs were brought forward for prioritization:

- Aging
- Children and Youth
- Safety
- Health Care Access
- Physical Health
- Mental Health

Members of the community stakeholder group determined that physical health and mental health services are the top unmet needs and priorities.

Sanford has determined the 2016-2019 implementation strategies for the following needs:

- Mental Health
- Physical Health

Implementation Strategies

Priority 1: Physical Health in the Community

- Obesity and inactivity are closely linked. Keeping active can help people stay at a healthy weight or lose weight. It can also lower the risk of heart disease, diabetes, stroke, high blood pressure, osteoporosis, and certain cancers, as well as reduce stress and boost mood. Inactive (sedentary) lifestyles do just the opposite.
- The World Health Organization, the U.S. Dept. of Health and Human Services, and other authorities recommend that for good health, adults should get the equivalent of two and a half hours of moderate-to-vigorous physical activity each week. Children should get even more, at least one hour a day.
- Sanford has made physical health a high priority, and has set measurable outcomes. The outcomes include assessing a baseline for weight and scheduled periodic measurement, education session participation and exercise participation. The Sanford Health *fit* program is available to all classroom teachers and includes four areas of focus; nutrition, activity, behavioral health and sleep. The program is available to promote healthy lifestyles in homes, schools, and daycares.

Priority 2: Depression

- Depression is a common but serious illness that can interfere with daily life. Many people with a depressive illness never seek treatment. But the majority, even those with the most severe depression, can get better with treatment. The North Dakota Department of Health reports that 14% of residents in Traill and Steele counties have reported fair or poor mental health days.
- Sanford has prioritized depression as a high priority and has set strategy to perform assessments for depression in the primary care setting and to improve PHQ-9 scores for patients who are diagnosed with depression. The goal is to improve PHQ-9 scores for patients with depression. The measurable outcome is the percentage of patients with major depression or dysthymia and an initial PHQ-9 score greater than nine whose six-month PHQ-9 score is less than five. Sanford Health will continue to provide a directory of services for community members.

Sanford Hillsboro Medical Center
Community Health Needs Assessment
2016

Table of Contents

| | Page |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Purpose of the Community Health Needs Assessment | 12 |
| Acknowledgements | 12 |
| Description of Sanford Hillsboro Medical Center | 15 |
| Description of the Hillsboro Community | 16 |
| Study Design and Methodology | 17 |
| Limitations of the Study | 18 |
| Key Findings | 19 |
| <ul style="list-style-type: none">• Community Health Concerns• Personal Health Concerns• Demographics• Health Needs and Community Resources Identified• Prioritization | |
| Addressing the Needs | 44 |
| 2016-2019 Implementation Strategies | 46 |
| 2013 Implementation Strategies Impact | 50 |
| Community Feedback from 2013 Community Health Needs Assessment | 53 |
| <u>Appendix</u> | 55 |
| <i>Primary Research</i> | 58 |
| <ul style="list-style-type: none">• <i>Asset Map</i>• <i>Prioritization Worksheet</i>• <i>Non-Generalizable Survey Results</i> | 62 |
| <i>Secondary Research</i> | 98 |
| <ul style="list-style-type: none">• <i>Trail County Community Health Profile</i>• <i>Steele County Community Health Profile</i>• <i>Definitions of Key Indicators / County Health Rankings</i>• <i>Trail County and Steele County Comparison</i> | |

Sanford Hillsboro Medical Center

Community Health Needs Assessment

2016

Purpose of the Community Health Needs Assessment

A community health needs assessment is critical to a vital community Benefit program that builds on community assets, promotes collaboration, improves community health, and promotes innovation and research. A community health needs assessment helps the community build capacity to support policy, systems, environmental changes, and community health improvement. A community health needs assessment also serves to validate progress made toward organizational strategies and provides further evidence for retaining not-for-profit status.

The purpose of this community health needs assessment is to develop a global view of the population's health and the prevalence of disease and health issues within our community. Findings from the assessment serve as a catalyst to align expertise and to develop a community benefit plan of action. There is great intrinsic value in a community health needs assessment when it serves to validate, justify and defend not-for-profit status and create opportunity to identify and address public health issues from a broad perspective.

Our guiding principles:

- All health care is a community asset
- Care should be delivered as close to home as possible
- Access to health care must be provided regionally
- Integrated care delivers the best quality and efficiency
- Community involvement and support is essential to success
- Sanford Health is invited into the communities we serve

Acknowledgements

Sanford Health would like to acknowledge and thank the steering committees for their assistance and expertise while performing the assessment and analysis of the community health data. The assessment provides support for the future direction of our work.

Sanford Enterprise Steering Group:

- JoAnn Kunkel, CFO, Sanford Enterprise
- Michelle Bruhn, CFO, Health Services Division
- Tiffany Lawrence, CPA, Fargo Region Co-Lead, CFO, Sanford Medical Center Fargo
- Bruce Viessman, Sioux Falls Region Co-Lead, CFO, Sanford Health Network
- Martha Leclerc, MS, Vice President, Revenue Management
- Randy Bury, CAO, Health Services Division
- Steve Goetsch, CFO, Sioux Falls Region
- Jane Heilmann, Senior Corporate Communications Strategist
- Joy Johnson, COO Bemidji Region
- Kirk Christy, CFO, Bismarck Region
- Carrie McLeod, MBA, MS, LRD,CDE, Enterprise Lead, Enterprise Community Health/Community Benefit

Sanford Hillsboro/Mayville Steering Group:

- Carrie McLeod, MBA, MS, LRD, CDE, Enterprise Lead, Community Health/Community Benefit
- Roger Baier, CEO- Sanford Mayville Medical Center
- Jac McTaggart, CEO, Sanford Hillsboro Medical Center

We express our gratitude to the following community collaborative members from across the enterprise footprint for their expertise with the planning, development and analysis of the community health needs assessment.

- Alicia Collura, Sioux Falls Public Health
- Anita Cardinal, Pennington County Public Health
- Ann Malmberg, Essentia Health
- Becky Secore, Beltrami Public Health
- Brenda Stallman, Traill County Public Health
- Brie Taralson, Essentia Health
- Brittany Ness, Steele County Public Health
- Caitlin Hurley, Avera Health
- Carrie McLeod, Sanford Health
- Dan Heinemann, MD, CMO, Sanford Health Network, Sioux Falls Region
- Gina Nolte, Partnership4Health, Clay County
- Hannah Shirkey, Sanford Health
- Jac McTaggart, Sanford Health
- Jessica Spaeth, City of Halstad
- Joy Johnson, Sanford Health
- Julie Jeske, CHI-St. Alexius Health
- Julie Miller, South Dakota Department of Health
- Julie Ward, Avera Health
- Kathy McKay, Clay County Public Health
- Katie Olson, South Dakota State University
- Kay Schwarzwalter, Center for Social Research, North Dakota State University
- Kim Jacobson, Traill County Public Health
- Kip Littau, South Dakota State University
- Marnie Walth, Sanford Health
- Mary Michaels, Sioux Falls Public Health
- Nancy Fahrenwald, South Dakota State University
- Renae Moch, Burleigh County Public Health
- Roger Baier, Sanford Health
- Ruth Bachmeier, Fargo Cass Public Health
- Sandra Melstad, Public Health Consultant, South Dakota Department of Health
- Stan Kogan, Sioux Falls Public Health
- Stephen Pickard, PhD, North Dakota Department of Health
- Susan Kahler, Burleigh County Public Health
- Teresa Miller, Avera Health

We extend special thanks to the representatives for the mentally and physically disabled, social services, non-profit organizations, and public health officers for their participation in this work. Together we are

reaching our vision “to improve the human condition through exceptional care, innovation and discovery”.

The following Mayville-Hillsboro-Halstad, Traill and Steele County key community stakeholders participated in community discussions and helped to formulate the priorities for future work:

- Jill Borth, Steele County Board of Health
- Deanna Killo, Mayville State University CDP Health
- Brittany Ness, Steele County Public Health
- Bonnie Overby, Mayville State University
- Casandra Garza, Mayville State University
- Brenda Stallman, Traill District Health
- Jessica Speath, Community of Halstad
- Ernie Strube, MayPort Economic Development Corporation
- Bill Krivarchka, Eastern North Dakota AHEC
- Jeff McInnes, Board Member, Sanford Hillsboro Medical Center
- JAc McTaggart, Sanford Hillsboro Medical Center
- Roger Baier, Sanford Medical Center



Description of Sanford Hillsboro Medical Center

Sanford Hillsboro consists of a 16-bed Critical Access Hospital, two provider-based Rural Health Clinics, a 36-bed long term care facility, and 16 assisted living units. The medical center employs approximately 130 people, including two family medicine physicians and a family medicine physician assistant. Services available at Sanford Hillsboro include 24/7 emergency care, general acute care, physical, occupational and speech therapy, radiology, IV therapy, lymphedema management, nutrition consultation, respite care, bariatric and pathology and FM Ambulance services.



Description of the Hillsboro Community

Hillsboro is a city of 1,600 people located on the banks of the Goose River in the heart of the Red River Valley of eastern North Dakota. It serves as the county seat for Traill County. Hillsboro was founded in 1869 by German settlers and named for railroad baron James J. Hill who stopped in the area in the late 1870s and found hospitality from a homesteader.

Hillsboro is conveniently located halfway between the metropolitan cities of Grand Forks and Fargo on Interstate 29. The surrounding area is prime farm land, and agriculture has dominated the area's economy from the beginning. One of the major industries and employers is American Crystal Sugar which operates a large sugar beet processing plant north of the city. The plant has been responsible for both an increase in population and a steady stream of available jobs.

Hillsboro has a strong economy with more than 60 businesses. It offers many amenities usually found in larger cities including K-12 education, medical facilities, parks, a pool, golf course and fitness center.

Study Design and Methodology

1. Non-Generalizable Survey

An on-line non-generalizable on-line survey was conducted through a partnership between the Sanford Hillsboro/Mayville/Halstad/Traill and Steel County Collaborative and the Center for Social Research (CSR) at North Dakota State University. The CSR developed and maintained links to the on-line survey tool. The website address for the survey instrument was distributed via e-mail to various agencies, at times using a snowball approach. Data collection occurred throughout the month of April 2015 and a total of 54 respondents participated in the on-line survey.

The purpose of this non-generalizable survey of community leaders was to learn about the perceptions of area community leaders regarding community health, their personal health, preventive health, and the prevalence of disease. This group included community leaders, legislators, and agency leaders representing chronic disease and disparity.

A Likert scale was developed to determine the respondent's highest concerns. Needs ranking 3.5 and above were included in the needs to be addressed and prioritized. As stated in the generalizable survey methodology, many of the identified needs that ranked below 3.5 are being addressed by Sanford. However, 3.5 and above was used as a focus for the purpose of the required prioritization.

2. Community Stakeholder Meeting

Community stakeholders were invited to a meeting to review the early findings from the generalizable survey and to discuss the top health issues or health-related issues facing the community. Community stakeholders helped to determine key priorities for the community, and these priorities are the focus of the implementation strategies for 2017-2019.

3. Community Asset Mapping

Asset mapping was conducted by reviewing the data and identifying the unmet needs from the various surveys and data sets. The process implemented in this work was based on the McKnight Foundation model - Mapping Community Capacity by John L. McKnight and John P. Kretzmann, Institute for Policy Research at Northwestern University.

Each unmet need was researched to determine what resources were available in the community to address the needs. The community stakeholder group conducted an informal gap analysis to determine what needs remained after resources were thoroughly researched. Once gaps were determined the group proceeded to the prioritization process. The multi-voting methodology was implemented to determine what top priorities would be further developed into implementation strategies.

4. Secondary Research

The secondary data includes community health profiles from the North Dakota Department of Health for Traill and Steele County, North Dakota. The Robert Wood Johnson County Health Rankings were also researched for Traill and Steel County.

Limitations of the Study

The findings in this study provide an overall snapshot of behaviors, attitudes, and perceptions of residents living in the Traill and Steele Counties in North Dakota and the area of Halstad, Minnesota. However, when comparing certain demographic characteristics (i.e., age, income, minority status) with the current population estimates from the U.S. Census Bureau, it was evident that white, more highly educated, and higher income earners were overrepresented. Overrepresentation of this nature is typical in health needs assessments.

A good faith effort was made to secure input from a broad base of the community. Invitations were extended to county and community leaders, organizations and agencies representing diverse populations and disparities.

The Internal Revenue Code 501 (r) statute requires that a broad base of key community stakeholders have input into the needs of the community. Those community members specified in the statute include: persons who represent the broad interests of the community served by the hospital facility including those with special expertise in public health; Federal, tribal, regional, state and or local health or other departments or agencies with information relevant to the health needs of the community served; leaders, representatives, or members of medically underserved, low-income, and minority populations.

Sanford extended a good faith effort to engage all of the aforementioned community representatives in the survey process. In some cases there were surveys that were submitted without names or without a specified area of expertise or affiliation. We worked closely with public health experts throughout the assessment process.

Public comments and response to the community health needs assessment and the implementations strategies are welcome on the Sanford website under “About Sanford” in the Community Health Needs Assessment section.


Key Findings

Community Health Concerns

Aging Population

The greatest area of concern among respondents is for the aging population, including the cost of long term care and the availability of memory care. Secondary research finds that 58.4 percent of the population in Traill County and 49.5 percent of the population in Steele County are over 65 years of age.

Level of concern with statements about the community regarding the AGING POPULATION




The North Dakota Long Term Care Association states that a primary reason for the concern about the cost of long term care is because people are not purchasing long term care insurance. When long term care services are required it may be a surprise to learn the true cost of the service.

Resources for family/friends and resources for the elderly to help them stay in their homes are becoming widely available through programs such as the North Dakota State University Extension Office and the Sanford Faith Community Nurse Program.

Children and Youth

The survey respondents have very high concerns for the children and youth of the community. The availability and cost of activities for children and youth, the availability and cost of quality child care, bullying, the availability and cost of quality infant care, and the availability and cost of services for at-risk youth are high concerns.


Level of concern with statements about the community regarding CHILDREN AND YOUTH



Health Care Access and Cost

Top priorities among survey respondents are access to affordable health insurance, the cost of affordable vision insurance, access to affordable prescription drug insurance, and the cost of affordable dental insurance.

Level of concern with statements about the community regarding HEALTH CARE



Access to care includes the ability to gain entry into a health system or provider service. Access can include the availability of health care providers and a workforce available to address the needs. Limited access can challenge the ability to receive appropriate levels of care and may pave the way to the utilization of higher cost entry points into the system through the emergency room.

The cost of care may influence the health of individuals. A Dartmouth study finds that North Dakota is considered a high quality state for health care in comparison to other states. North Dakota is also considered to deliver health care at a low cost in comparison to other states.

Sanford provides a community care program and financial assistance policy to address assistance to all who qualify for charity care. During fiscal year 2014 Sanford contributed over \$51 million for charity care for our patient population that required care without the ability to pay for services. Sanford has financial counselors available at all clinic and medical center facilities to assist patients with applications for assistance and access needs.

One example of a community resource that is addressing the access needs of patients is Sanford's My Sanford Nurse program (formerly called Ask-A-Nurse). My Sanford Nurse served 324,295 individuals from throughout the footprint and nation during fiscal year 2014 and provided a community benefit of over \$1.8M with more than 45,965 nursing staff hours. There is no fee for this service.

Timely access for mental health and behavioral health providers continues to remain a concern across the community. Workforce gaps among mental health and behavioral health providers remains a challenge. Sanford has an aggressive recruiting program and continues to recruit these experts to our community. The North Dakota Human Services Behavioral Health Interim Committee is also focused on developing the behavioral health workforce for the state in the areas of adult psychiatry, child and adolescent psychiatry, and substance abuse providers. Sanford leaders are serving on the stakeholder group with this interim committee, along with other community providers.

Physical and Mental Health


Survey respondents have high levels of concern for physical and mental health issues. The top concerns are cancer and chronic disease. Additional high ranking concerns include inactivity and lack of exercise, obesity, and poor nutrition. Stress and depression rank the highest concerns for mental health.

Secondary research through the 2015 County Health Rankings find that the average number of self-reported mentally unhealthy days reported in the last 30 days is at 2.2 days in Traill County and 2.3 days in Steele County.

Traill County residents reporting poor or fair health is at 13% and Traill County is at 12%. Traill County residents reporting that they are obese are at 36%; Traill County is at 32%. Traill County residents who report being physically inactive is at 27%; 30% of the residents in Steele County report being physically inactive. (See Appendix)

The North Dakota Department of Health reports that 9% of Traill and Steele County residents have been told by their health provider that they have cancer (excluding skin cancer). (See Appendix)

Level of concern with statements about the community regarding PHYSICAL AND MENTAL



Concerns about cancer: The prevalence of cancer among community members is on the minds of our residents. The Roger Maris Cancer Center in Fargo, North Dakota is located in close proximity to Mayville.

Cancer is the second leading cause of death in the United States for the adult population. According to the UND School of Medicine and Health Sciences 2015 Biennial Report, digestive system cancer is the most commonly diagnosed cancer in our state, followed by breast cancer. The study finds that the risk of cancer in North Dakota is higher than in the rest of the nation.

Sanford is committed to serving the cancer needs of our community through the Roger Maris Cancer Center and the 13 medical oncologists/hematologists, 5 radiation oncologists, 2 pediatric oncologists/hematologists, 7 palliative care board certified physicians, 1 oncology clinical psychologist, 1 medical geneticist, 3 genetic counselors, 1 doctor of nursing practice, 2 physician assistants, 2 nurse practitioners, 7 pharmacists, 4 radiation oncology medical physicists, and several hundred nurses.

Sanford's Roger Maris Cancer Center holds a number of accreditations including Joint Commission Accreditation, the American College of Surgeons Commission on Cancer – Silver Commendation, the National Accreditation Program for Breast Cancer through the American College of Surgeons Accreditation by the American College of Radiology – Radiation Oncology, Edith Sanford Breast Health Comprehensive Center recognized as a Certified Quality Breast Center of Excellence through the National Consortium of Breast Centers, Breast Imaging Center of Excellence through the American College of Radiology, Certification by the Quality Oncology Practice Initiative through the American Society of Clinical Oncology, and the Advanced Care Certification by The Joint Commission for Palliative Care Services.

The Sanford Roger Maris Cancer Center desires to detect cancer at earlier stages when there is a better chance for cure. One example of the Center's focus on early detection is colorectal screenings. Colorectal cancer screenings can detect precancerous colorectal polyps which, if removed, can prevent colorectal cancer and reduce an individual's colorectal cancer risk by 50%.


In 2014 at Sanford Roger Maris Cancer Center there were 158 colorectal cancer cases, 74 (47%) of which were more advanced (stage 3 or 4). Treatment results compared favorably with other cancer centers in the region and nationally. North Dakota is at the bottom third of the country for colorectal cancer screenings, which leads to later detection and more advanced colorectal cancer stages.

Having one of the familial cancer syndromes increases the likelihood of developing cancer. The Roger Maris Cancer Center has a familial cancer syndrome clinic, where the team assists in coordinating care and helping individuals with an increased risk of cancer to plan for future screenings, and customizes the patient's treatment plan. Genetic risk assessment is an important component of the care at the Sanford Roger Maris Cancer Center.

Concerns about other chronic diseases: The top chronic diseases among respondents include hypercholesterolemia, hypertension, arthritis, and diabetes. Respondents were also concerned about comorbidities such as obesity, poor nutrition and lack of physical activity.

The North Dakota Department of Health reports that 24.2% of Traill and Steele County residents have been diagnosed with arthritis. The state health data also reports that 76.2% of Traill and Steele County residents are overweight or obese, which makes it more uncomfortable when dealing with arthritis. Traill and Steele County residents who report that they have been told by their provider that they have hypertension is 24%, and 40.3% report the same for high cholesterol. (See Appendix)

Whether respondents have any of the following chronic diseases



Concerns about hypertension: Hypertension is addressed at Sanford through a standardized protocol, frequent blood pressure monitoring, and referral throughout all departments. Continued improvement has been realized since the implementation of the standardized protocol.

Concerns about hypercholesterolemia: Sanford providers address hypercholesterolemia through evidence-based guidelines. Patients are closely monitored to assure a reduction in heart disease risk.

Concerns about diabetes: Sanford offers a comprehensive diabetes education program. Sanford diabetes clinics and centers are dedicated to empowering people with diabetes to feel better and prevent long-term complications. Sanford offers assessment and personalized education care to give patients and their families the tools they need to manage diabetes while living well. Endocrinologists, certified diabetes nurses and certified diabetes dieticians provide diagnosis, assessment, one-on-one education, and instruction.

The chronic disease self-management Better Choices, Better Health program at Sanford is offered free of charge to community members. Better Choices, Better Health is modeled after Stanford University's chronic disease self-management program. The workshops are 2½ hours long and meet weekly for 6 weeks. The program is facilitated by two trained lay leaders and one or both have a chronic condition. Research has found that after participating in the program individuals are better able to manage their symptoms, communicate more easily with their doctors, are less limited by the disease, and generally feel better.

Concerns about obesity, poor nutrition and lack of physical activity: Sanford Enterprise chose obesity as an implementation strategy for the 2013 CHNA. Strategies included an annual obesity symposium for providers and monthly education programs for community members. The Sanford obesity symposium was attended by more than 400 registrants during April 2013 and March 2015. The symposium will be held again during March 2016. The symposium is evidence-based. Sanford providers and national experts serve as faculty for the symposium.

The Sanford Health *fit* initiative, <http://sanfordfit.org/> a childhood obesity prevention initiative, continues to grow and mature as we work to refine the offerings and enable broad replication and meaningful use. Supported by the clinical experts of Sanford Health, *fit* educates, empowers and motivates families to live a healthy lifestyle through a comprehensive suite of resources for children, parents, teachers and clinicians. *fit* is the only initiative focusing equally on the four key contributing factors to childhood obesity: Food (nutrition), Move (activity), Mood (behavioral health), and Recharge (sleep). Sanford's *fit* Initiative has come a long way since its inception in 2010. Through *fit* we are actively working to promote healthy lifestyles in homes, schools, daycares, our clinical settings, and throughout the community by way of technology, engaging programs, and utilizing key role models in a child's life.

- The *fit* website for Juniors, Kids and Teens creates an entertaining and interactive on-line environment where they can play games, watch videos and take daily challenges. Parents benefit from their own set of resources where they can find tips and tools on becoming healthy role models and raising *fit* kids. To date, the children's and parent's sites have received more than 7.5 million visitors. Over 700 pieces of content have been added to the sites including videos, slideshows, games, articles, and even *fit* songs.
- In addition to the web, *fit* is developing meaningful school resources to bring value and fun into the primary education setting. We are doing this by integrating *fit* points into science and math components to provide health promotion, an avenue into the classroom without taking valuable time away from those critical subjects.
- *fit4Schools* – *fit4Schools* includes unique *fit*-based lessons integrated into daily classroom activities. fit4schools.sanfordfit.org is an on-line school resource that incorporates topics into math and science curriculum. To date the program has 14 STEM (integrating science, technology, engineering, and math) unit plans that can be downloaded for classroom use. To date it has:
 - Reached 50,000 schools
 - 180,000 page views from educators across the country
 - 12,000 lesson plan downloads, representing 600,000+ students
- We are also reaching thousands of students through several pilot school programs.
 - *fit4Schools* – *fit4Schools*, which includes unique *fit*-based lessons integrated into daily classroom activities, is in its final phase of development. It is being piloted in seven elementary schools in the Sanford region.
- Community
 - The *fit* friends, Denny, Abby, Sam, Alex and Marty, along with the *fit* team, have been making a variety of appearances at events across the Sanford footprint. *fit* has been at over 2 dozen events interacting with more than 15,000 children and parents to spread the word about the *fit* platform and resources.
 - Smartphone Apps – Through a series of fun and engaging apps, *fit* will continue to activate kids at the touch of a fingertip to live a fit and healthy lifestyle related to Mood, Recharge, Food and Move.
 - MOVE2Draw is a simple and fun way for kids to move and create their own unique drawings. Once a drawing is completed, it can be stored on the MOVE2Draw website.

- eMOODicam is a photo application that allows the user to enhance a photo and bring the mood to life and share with others.
- Looking Forward
 - *fit* is continuing to look to the future for ways to continue to make a meaningful impact on children and families both on-line and off-line. Other exciting expansions that are in the works include:
 - Clinical Setting – Resources for the clinical setting to spur actionable and understandable discussions between health care providers and families.
 - Health Coaches – Exploring meaningful ways for health coaches to promote healthy choices with children and adults.
 - Engage Key Role Models – Firefighters and youth sport coaches are role models and have a big influence on children so that's why *fit* is developing resources for them to teach the principles of *fit* along with sports fundamentals and other outreach efforts.
 - *fit*Club 4 Boys – 10-week after school program for boys, ages 8-12, to develop knowledge of *fit* principles and healthy behavior choices.
 - *fit* Parent/child – Class for parents and children to understand healthy choices and the benefits of living a healthy lifestyle.



Concerns about mental/behavioral health: During the 2013 CHNA process, Sanford Health developed an implementation strategy to fully integrate behavioral health services or access to behavioral health outreach in all regional clinic sites in the Fargo, Sioux Falls, Bismarck and Bemidji regions.


A key implementation strategy for Sanford Health is to fully integrate behavioral health services into all primary care clinics. Sanford One Care is addressing behavioral health in primary and specialty care clinics and medical centers. Sanford One Care uses technology to conduct behavioral health screenings in order to identify behavioral health concerns as early as possible. Through deployment of Medical Home with fully integrated behavioral health care services, Sanford is providing patient-centered care collaborative teams to meet the needs of Sanford patients.

- During 2015 Sanford hosted a behavioral health symposium. The symposium was designed to enhance the competence of health care professionals in the identification, treatment and management of behavioral health issues impacting specialty and primary care.

Sanford behavioral health experts are serving on the North Dakota Behavioral Health Stakeholders Advisory to the Department of Human Services Legislative Interim Committee to shape policy. The North Dakota Behavioral Health Stakeholders group works to address unmet needs in behavioral health in North Dakota. Using a Substance Abuse and Mental Health Services Administration (SAMHSA) template of the components of a comprehensive system of behavioral health care, the group has identified needs in North Dakota and is working on solutions to meet those needs. Types of behavioral health services addressed by this group include adult mental health, children's mental health, substance abuse, and workforce development.

Concerns about substance use and abuse: Survey respondents have a high level of concern about alcohol use and abuse in the community and about underage drinking and use of drugs. Additionally, smoking and tobacco use and drug use and abuse among adults are high concerns.

Level of concern with statements about the community regarding SUBSTANCE USE AND ABUSE




Safety

The highest concerns for survey respondents in the safety category are the presence of drugs and alcohol in the community, child abuse and neglect, and domestic abuse.

Secondary research finds that alcohol-impaired driving deaths have reached 33% in Traill County and 50% in Steele County. (See Appendix)

Level of concern with statements about the community regarding SAFETY




Sanford is participating in the North Dakota “Reducing Pharmaceutical Narcotics in Our Communities - Through Education and Awareness” committee. The committee has a four-pillar approach including: education and awareness, prescription drug-take back program, law enforcement, pharmacy partnership, and the prescription drug monitoring program.

Personal Health Concerns

Respondents' Personal Health Status


The study results suggest possible discrepancies between respondents' perceived personal health and their actual health status as determined by objective measures. For example, using the Body Mass Index (BMI) which calculates weight status using an individual's weight and height, the majority of respondents are overweight or obese. However, the vast majority of community respondents rate their own health as excellent, very good or good.

Respondents' rating of their health in general




- 88.7% of the respondents rate their health as good or better

Respondents' weight status based on the Body Mass Index (BMI) scale



- 65.3% of respondents report a BMI that is overweight or obese

Length of time since respondents last visited a doctor or health care provider for a routine physical exam and length of time since they last visited a dentist or dental clinic for any reason



- 21% of respondents have waited longer than a year to visit a doctor or health care provider
- 17.3 % of respondents have waited more than a year to see a dentist

Preventive Health

Preventive health care promotes the detection and prevention of illness and disease and is another important component of good health and well-being. Community results indicate that within the past year, the majority of respondents had a blood pressure screening, blood sugar screening, cholesterol screening, dental screening, flu shot, pelvic exam (females), breast cancer screening (females), and cervical cancer screening (females). However, there are many screenings and tests that a majority of respondents did not receive (i.e., bone density test, cardio screening, glaucoma test, hearing screening, immunizations, STD test, vascular screening, colorectal cancer screening, prostate cancer screening (males), and skin cancer screening in the past year. Many tests and screenings may be conditional upon guidelines, which can be age sensitive/appropriate.

Whether or not respondents have had preventive screenings in the past year, by type of screening

| Type of screening | Yes | No |
|-----------------------------------------|-------------|-------------|
| GENERAL SCREENINGS | | |
| Blood pressure screening (N=380) | 86.5 | 13.5 |
| Blood sugar screening (N=379) | 64.7 | 35.3 |
| Bone density test (N=367) | 6.0 | 94.0 |
| Cardiovascular screening (N=373) | 24.0 | 76.0 |
| Cholesterol screening (N=378) | 67.3 | 32.7 |

| Type of screening | Yes | No |
|---------------------------------------------------|------|------|
| Dental screening and X-rays (N=378) | 82.7 | 17.3 |
| Flu shot (N=380) | 76.9 | 23.1 |
| Glaucoma test (N=374) | 50 | 50 |
| Hearing screening (N=377) | 6.1 | 93.9 |
| Immunizations (tetanus, hepatitis A or B) (N=374) | 22 | 78 |
| Pelvic exam (N=189 Females) | 60 | 40 |
| STD (N=369) | 12 | 88 |
| Vascular screening (N=368) | 11.8 | 88.2 |
| CANCER SCREENINGS | | |
| Breast cancer screening (N= 189 Females) | 75.6 | 24.4 |
| Cervical cancer screening (N=185 Females) | 60 | 40 |
| Colorectal cancer screening (N=368) | 18 | 82 |
| Prostate cancer screening (N=182 Males) | 0 | 100 |
| Skin cancer screening (N=365) | 14.3 | 85.7 |

- For most types of screenings, the most common reasons for not getting the test or procedure are that it is not necessary and the doctor has not suggested one
- For the flu shot screening, most respondents cite other reasons for not getting the shot
- 44% of the survey respondents were under 45 years of age

Screenings

Breast cancer screening: According to the Center for Disease Control (CDC), a mammogram is an X-ray of the breast. Mammograms are the best way to find breast cancer early, when it is easier to treat and before it is big enough to feel or cause symptoms. Having regular mammograms can lower the risk of dying from breast cancer. The United States Preventive Services Task Force recommends that if you are 50 to 74 years old, be sure to have a screening mammogram every two years. If you are 40 to 49 years old, talk to your doctor about when to start and how often to get a screening mammogram.

Cervical cancer screening: Cervical cancer is the easiest gynecologic cancer to prevent, with regular screening tests and follow-up. Two screening tests can help prevent cervical cancer or find it early:

- The Pap test (or Pap smear) looks for *pre-cancers*, cell changes on the cervix that might become cervical cancer if they are not treated appropriately.
- The HPV test looks for the virus that can cause these cell changes (human papillomavirus) (http://www.cdc.gov/cancer/hpv/basic_info/)
- The Pap test is recommended for all women between the ages of 21 and 65 years old, and can be done in a doctor's office or clinic.

Colorectal cancer screening: Colorectal cancer almost always develops from *precancerous polyps* (abnormal growths) in the colon or rectum. Screening tests can also find colorectal cancer early, when treatment works best. Regular screening, beginning at age 50, is the key to preventing colorectal cancer.

The U.S. Preventive Services Task Force (USPSTF) recommends screening for colorectal cancer using high-sensitivity fecal occult blood testing, sigmoidoscopy, or colonoscopy beginning at age 50 and continuing until age 75.

Prostate cancer screening: The American Cancer Society (ACS) recommends that men have a chance to make an informed decision with their health care provider about whether to be screened for prostate cancer. The decision should be made after getting information about the uncertainties, risks, and potential benefits of prostate cancer screening. Men should not be screened unless they have received this information. The discussion about screening should take place at:

- Age 50 for men who are at average risk of prostate cancer and are expected to live at least 10 more years.
- Age 45 for men at high risk of developing prostate cancer. This includes African Americans and men who have a first-degree relative (father, brother or son) diagnosed with prostate cancer at an early age (younger than age 65).
- Age 40 for men at even higher risk (those with more than one first-degree relative who had prostate cancer at an early age).

After this discussion, those men who want to be screened should be tested with the prostate-specific antigen (PSA) blood test. The digital rectal exam (DRE) may also be done as a part of screening.

If, after this discussion, a man is unable to decide if testing is right for him, the screening decision can be made by the health care provider, who should take into account the patient's general health preferences and values.

Assuming no prostate cancer is found as a result of screening, the time between future screenings depends on the results of the PSA blood test:

- Men who choose to be tested who have a PSA of less than 2.5 ng/mL may only need to be retested every 2 years.
- Screening should be done yearly for men whose PSA level is 2.5 ng/mL or higher.

Because prostate cancer often grows slowly, men without symptoms of prostate cancer who do not have a 10-year life expectancy should not be offered testing since they are not likely to benefit. Overall health status, and not age alone, is important when making decisions about screening.

Even after a decision about testing has been made, the discussion about the pros and cons of testing should be repeated as new information about the benefits and risks of testing becomes available. Further discussions are also needed to take into account changes in the patient's health, values and preferences.

Skin cancer screening: The U.S. Preventive Services Task Force (USPSTF) has concluded there is not enough evidence to recommend for or against routine screening (total body examination by a doctor) to find skin cancers early. The USPSTF recommends that doctors:

- Be aware that fair-skinned men and women aged 65 and older, and people with atypical moles or more than 50 moles, are at greater risk for melanoma.
- Look for skin abnormalities when performing physical examinations for other reasons.

Flu Vaccines

The Center for Disease Control's Advisory Committee on Immunization Practices (ACIP) recommends that everyone six months and older receive a flu vaccine annually. Findings from the generalizable survey indicate that 37% of respondents did not have a flu shot last year.

The Center for Disease Control states that influenza is a serious disease that can lead to hospitalization and sometimes even death. Even healthy people can get sick from the flu and spread it to others. Flu vaccines cause antibodies to develop in the body about two weeks after vaccination. These antibodies provide protection against infection with the viruses that are in the vaccine.


Fruit and Vegetable Intake

The study results suggest that the majority of respondents do not meet vegetable and fruit recommended dietary guidelines. Only 28.4% of respondents reported having 3 or more servings of vegetables the prior day, and 17% of the group reported having 3 or more servings of fruits the prior day.

According to the U.S. Department of Health and Human Services, U.S. Department of Agriculture Dietary Guidelines for Americans, it is recommended that individuals consume 3 to 5 servings of vegetables per day and 2 to 4 servings of fruit per day depending on age. A diet high in fruits and vegetables is associated with decreased risk for chronic diseases. In addition, because fruits and vegetables have low energy density (i.e., few calories relative to volume), eating them as part of a reduced-calorie diet can be beneficial for weight management.

Secondary research provided by the North Dakota Department of Health finds that 84.4% of residents in Traill and Steele County report not getting 5 fruits and vegetables each day. (Appendix page__)

Number of servings of vegetables, fruit, and fruit juice that respondents had yesterday




Physical Activity Levels

Study results suggest that the majority of respondents do not meet physical activity guidelines. 39.3% of respondents engage in moderate activity 3 or more times per week and 17% engage in vigorous activity 3 or more times per week.

Guidelines from the Centers for Disease Control and Prevention recommend that individuals participate in 150 minutes of moderate physical activity per week or 75 minutes of vigorous physical activity per week to help sustain and improve health.

Number of days in an average week respondents engage in MODERATE and VIGOROUS activity




Tobacco Use


Study results indicate that the vast majority of community respondents are not currently tobacco users. However, over 37% of respondents have smoked at least 100 cigarettes in their lifetime, which indicates former smoker status according to the Centers for Disease Control and Prevention.

Secondary research through the 2015 County Health Rankings finds that 13% of Traill County residents and 12% of Steele County residents are current smokers.

Whether respondents have smoked at least 100 cigarettes in their entire life



How often respondents currently smoke cigarettes and use chewing tobacco or snuff




Mental Health


Mental health is an important component of well-being at every stage of life and impacts how we think, act and feel. Mental health influences our physical health, how we handle stress, how we make choices, and how we relate to others. Among survey respondents, mental health is a

moderately high area of concern, particularly depression and stress. 37% of respondents have been told or diagnosed by a doctor or health professional that they have anxiety or stress and 31.5% have been told they have depression. In addition, more than half of respondents self-report that in the last month, there were days when their mental health was not good. 55.8% of the respondents reported days in the past month when their mental health was not good.

Percentage of respondents who have been told by a doctor or health professional that they have a mental health issue, by type of mental health issue



Number of days in the last month that respondents' mental health was not good




Substance Abuse Responses


Substance abuse is also a mental health disorder, as defined by the *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition (DSM-IV), and can stem from mental health concerns. In the community, 74% of respondents drank alcoholic beverages on at least one day in the last month. On days they drank, 29.7% of respondents drank an average of three or more drinks per day. In regards to binge drinking, 26.5% of respondents report binge drinking at least once per month.

Secondary research through the 2015 County Health Ranking indicates that 16% of Traill County and 26% of Steele County residents report binge drinking. (See Appendix)

Number of times during the past month that respondents consumed at least 4 or 5 alcoholic drinks (Binge drinking is defined by the CDC as 4 drinks for females, 5 drinks for males) on the same occasion




Whether respondents have ever had a problem with alcohol use or prescription or non-prescription drug abuse



Less than 5% of respondents reported having a problem with alcohol although earlier reporting indicated a higher level of binge drinking. Overall, 32.7% of respondents report alcohol use has had harmful effects on themselves or a family member.

Other forms of substance abuse include the use of prescription or non-prescription drugs. No respondents reported having had a problem with prescription or non-prescription drug abuse. However, 15.4% of respondents say prescription or non-prescription drug abuse has had harmful effects on themselves or a family member.

Whether alcohol use or prescription or non-prescription drug abuse has had harmful effects on respondents or a family member over the past two years



Demographics

Total Population – 2010 U.S. Census Bureau

- Traill County: 8,121
- Steele County: 1,975

Population by Age and Gender

| | Number | Percent | Males | Percent | Females | Percent |
|--------------------|------------------------------|----------------------------|----------------------------|----------------------------|------------------------------|----------------------------|
| <5 years | Traill: 510 Steele: 101 | Traill: 6.3 Steele: 5.1 | Traill: 256 Steele: 45 | Traill: 3.2 Steele:2.3 | Traill: 254 Steele: 56 | Traill: 3.1 Steele: 2.8 |
| 5-9 | Traill: 465 Steele: 94 | Traill: 5.7 Steele: 4.8 | Traill: 238 Steele: 46 | Traill: 2.9 Steele: 2.3 | Traill: 227 Steele: 48 | Traill: 2.8 Steele: 2.4 |
| 10-14 | Traill: 504 Steele: 134 | Traill: 6.2 Steele: 6.8 | Traill: 268 Steele: 58 | Traill: 3.3 Steele: 2.9 | Traill: 236 Steele: 76 | Traill: 2.9 Steele: 3.8 |
| 15-19 | Traill: 604 Steele: 121 | Traill: 7.4 Steele: 6.1 | Traill: 330 Steele: 60 | Traill: 4.1 Steele: 3.0 | Traill: 274 Steele:61 | Traill: 3.4 Steele: 3.1 |
| 20-24 | Traill: 528 Steele: 62 | Traill: 6.5 Steele: 3.1 | Traill: 286 Steele: 31 | Traill: 3.5 Steele: 1.6 | Traill: 242 Steele: 31 | Traill: 3.0 Steele: 1.6 |
| 25-29 | Traill: 385 Steele: 95 | Traill: 4.7 Steele: 4.8 | Traill: 204 Steele: 57 | Traill: 2.5 Steele: 2.9 | Traill: 181 Steele: 38 | Traill: 2.2 Steele: 1.9 |
| 30-34 | Traill: 401 Steele: 90 | Traill: 4.9 Steele: 4.6 | Traill: 207 Steele: 52 | Traill: 2.5 Steele: 2.6 | Traill: 194 Steele: 38 | Traill: 2.4 Steele: 1.9 |
| 35-39 | Traill: 428 Steele: 86 | Traill: 5.3 Steele: 4.4 | Traill: 214 Steele: 41 | Traill: 2.6 Steele: 2.1 | Traill: 214 Steele: 45 | Traill: 2.6 Steele: 2.3 |
| 40-44 | Traill: 482 Steele: 118 | Traill: 5.9 Steele: 6.0 | Traill: 232 Steele: 70 | Traill: 2.9 Steele: 3.5 | Traill: 250 Steele: 48 | Traill: 3.1 Steele: 2.4 |
| 45-49 | Traill: 594 Steele: 178 | Traill: 7.3 Steele: 9.0 | Traill: 328 Steele: 85 | Traill: 4.0 Steele: 4.3 | Traill: 266 Steele: 93 | Traill: 3.3 Steele: 4.7 |
| 50-54 | Traill: 643 Steele: 157 | Traill: 7.9 Steele: 7.9 | Traill: 321 Steele: 88 | Traill: 4.0 Steele: 4.5 | Traill: 322 Steele: 69 | Traill: 4.0 Steele: 3.5 |
| 55-59 | Traill: 604 Steele: 165 | Traill: 7.4 Steele: 8.4 | Traill: 321 Steele: 89 | Traill: 4.0 Steele:4.5 | Traill: 283 Steele: 76 | Traill: 3.5 Steele: 3.8 |
| 60-64 | Traill: 452 Steele: 133 | Traill: 5.6 Steele: 6.7 | Traill: 234 Steele: 80 | Traill: 2.9 Steele: 4.1 | Traill: 218 Steele: 53 | Traill: 2.7 Steele: 2.7 |
| 65-69 | Traill: 380 Steele: 113 | Traill: 4.7 Steele: 5.7 | Traill: 170 Steele: 58 | Traill: 2.1 Steele: 2.9 | Traill: 210 Steele:55 | Traill: 2.6 Steele: 2.8 |
| 70-74 | Traill: 282 Steele: 104 | Traill: 3.5 Steele: 5.3 | Traill: 135 Steele: 47 | Traill: 1.7 Steele: 2.4 | Traill: 147 Steele:57 | Traill: 1.8 Steele:2.9 |
| 75-79 | Traill: 260 Steele: 105 | Traill: 3.2 Steele: 5.3 | Traill: 103 Steele:48 | Traill: 1.3 Steele: 2.4 | Traill: 157 Steele:57 | Traill: 1.9 Steele: 2.9 |
| 80-84 | Traill: 279 Steele: 58 | Traill: 3.4 Steele: 2.9 | Traill: 124 Steele: 23 | Traill: 1.5 Steele: 1.2 | Traill: 155 Steele: 35 | Traill: 1.9 Steele: 1.8 |
| 85 and over | Traill: 320 Steele: 61 | Traill: 3.9 Steele: 3.1 | Traill: 96 Steele:31 | Traill: 1.2 Steele:1.6 | Traill: 224 Steele:30 | Traill: 2.8 Steele: 1.5 |
| | | | | | | |
| Median age | Traill: 42.6 Steele: 47.7 | | Traill: 40.9 Steele: 48 | | Traill: 44.3 Steele: 47.3 | |

Population by Race

| | Traill | Percent | Steele | Percent |
|-------------------------------------------|---------------|----------------|---------------|----------------|
| White | 7,809 | 96.2 | 1,927 | 92.7 |
| Black or African American | 42 | 0.5 | 3 | 0.2 |
| American Indian or Alaska Native | 64 | 0.8 | 23 | 1.2 |
| Asian | 21 | 0.3 | 2 | 0.1 |
| Native Hawaiian or other Pacific Islander | 1 | 0.0 | 0 | 0.0 |
| Hispanic or Latino | 214 | 2.6 | 20 | 1.0 |

The per capita personal income in Traill County, North Dakota is \$27,096. The per capita personal income in Steele County is \$31,455. Those living below the poverty level are 10.7% in Traill County and 6.2% in Steele County. The unemployment rate in Traill County is 3.1% and the unemployment level in Steele County is 2.2%.

Health Needs and Community Resources Identified

One of the requirements for a community health needs assessment is to identify the resources that are available in the community to address unmet needs. Asset mapping was conducted by reviewing the primary and secondary research and identifying the unmet needs from the various surveys and data sets. Each unmet need was researched to determine what resources were available in the community to address the needs.

The key community stakeholders from Hillsboro, Portland, Mayville, Halstad, Traill and Steel County participated in the asset mapping and reviewed the research findings. The group conducted an informal gap analysis to determine what needs remained after resources were thoroughly researched. Once gaps were determined the group proceeded to the prioritization process. The multi-voting methodology was implemented to determine what top priorities would be further developed into implementation strategies.

The process implemented in this work was based on the McKnight Foundation model - Mapping Community Capacity by John L. McKnight and John P. Kretzmann, Institute for Policy Research at Northwestern University.

The asset map includes identified needs from the following:

- Identified needs from the non-generalizable survey
- Concerns expressed by the key stakeholder group
- Secondary research data
- Community resources that are available to address the need(s)

The asset map can be found in the Appendix.

Prioritization

The following needs were brought forward for prioritization:

- Aging
- Children and Youth
- Safety
- Health Care Access
- Physical Health
- Mental Health
- Preventive Health

Sanford is addressing all of the assessed needs that fall within our scope of work. In some cases the need is one where we do not have the expertise to adequately address the need; however, Sanford leaders will communicate these findings to community leaders and experts who can best focus on a solution to the concern.

A document that shares what Sanford is doing to address the need or defends why Sanford is not addressing the need can be found in the Appendix.

Members of the community stakeholder group determined that mental health and physical health are the top unmet needs.

Sanford has determined the 2016-2019 implementation strategies for the following needs:

- Mental Health
- Physical Health

Addressing the Needs

2016 Community Health Needs Assessment Sanford Hillsboro

| Identified Concerns | How Sanford Hillsboro is Addressing the Needs |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>Aging</p> <ul style="list-style-type: none"> • Cost of long term care • Availability of memory care | <p>The cost of LTC is governed by the State of North Dakota. We can continue to control our costs but North Dakota Human Services determines the actual costs.</p> <p>When we built our LTC facility in 2008 a study was completed to show the need for memory care, the previous board decided that was not our business. Now the availability to purchase LTC beds for such a purpose is not an option, along with our facility has no property to expand and our LTC beds run full.</p> |
| <p>Children and Youth</p> <ul style="list-style-type: none"> • Bullying • Availability of activities for children and youth • Cost of activities for children and youth | <p>The local school has a program for bullying. As part of Physical Health, will be working with high school and local public health unit in determining where child activities are happening and what can be provided as we are a very small community.</p> |
| <p>Safety</p> <ul style="list-style-type: none"> • Seat belt usage • Presence of street drugs and alcohol in the community | <p>City council and County sheriff's office are working on promotion of seat belt usage and also Sheriff's Office is monitoring illegal drugs and alcohol consumption.</p> |
| <p>Health Care</p> <ul style="list-style-type: none"> • Access to affordable health insurance • Cost of affordable vision insurance • Access to affordable prescription drugs • Cost of affordable dental insurance coverage | <p>Have and will continue to have community forums on these topics presented by various Health Plans. We are only participants as we are not in business of selling insurance.</p> |
| <p>Physical Health</p> <ul style="list-style-type: none"> • Cancer • Poor nutrition and eating habits • Inactivity and lack of exercise • Obesity • Chronic disease • BMI – overweight or obese • Only 25.9% of respondents have 3 or more vegetables/day and 25.4 % have 3 or more fruits/day • Only 45.5% have 3 or more days each week of moderate activity and 20 % report 3 or more | <p>Sanford continues to promote wellness.</p> <p>The 2017-2019 implementation strategies are set to address physical health</p> <p>See Community Health Needs Assessment – Implementation Strategy for Sanford Hillsboro Medical Center for more in depth explanations.</p> |

| Identified Concerns | How Sanford Hillsboro is Addressing the Needs |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>days of vigorous activity each week</p> <ul style="list-style-type: none"> • 30.9% have smoked at least 100 cigarettes in their life • 20.3% of respondents reported hypertension • 16.9% reported high cholesterol • 10.2% reported diabetes | |
| <p>Mental Health</p> <ul style="list-style-type: none"> • Stress • Depression • Dementia and Alzheimer’s • 22% of respondents report that they have been told by a doctor that they have anxiety or stress, and 15.3% report being told that they have depression • 59.1% reported 1 or more days in the last month when their mental health was not good. • 11.9% of respondents reported 3 or more drinks /d on average • 26% reported 4 or 5 drinks (binge) on the same occasion over the past month • No one reported having a problem with alcohol use or drug use, however 24.1% reported that alcohol use had harmful effects on the respondent or a family member | <p>Continuing to update a directory for available resources in county.</p> <p>Have secured a small grant to go to 9 senior citizen centers in area to talk about physical exercise (includes diet) and chronic disease. We hope to have a health fair also to educate the public.</p> |
| <p>Preventive health</p> <ul style="list-style-type: none"> • Flu shots and immunizations | <p>Sanford conducts flu blitzes in the community, and when patients come into clinic, immunizations are recommended.</p> |

2016-2019 Implementation Strategies

Implementation Strategies

Priority 1: Physical Health in the Community

- Obesity and inactivity are closely linked. Keeping active can help people stay at a healthy weight or lose weight. It can also lower the risk of heart disease, diabetes, stroke, high blood pressure, osteoporosis, and certain cancers, as well as reduce stress and boost mood. Inactive (sedentary) lifestyles do just the opposite.
- The World Health Organization, the U.S. Dept. of Health and Human Services, and other authorities recommend that for good health, adults should get the equivalent of two and a half hours of moderate-to-vigorous physical activity each week. Children should get even more, at least one hour a day.
- Sanford has prioritized physical health and has set measurable outcomes that include assessing a baseline weight and periodic measurement, education session participation and exercise participation. The Sanford Health *fit* program is available to all classroom teachers and includes four areas of focus; nutrition, activity, behavioral health and sleep. The program is available to promote healthy lifestyles in homes, schools, and daycares.

Priority 2: Depression

- Depression is a common but serious illness that can interfere with daily life. Many people with a depressive illness never seek treatment. But the majority, even those with the most severe depression, can get better with treatment. The North Dakota Department of Health reports that 14% of residents in Traill and Steele counties have reported fair or poor mental health days.
- Sanford has prioritized depression as a top priority and has set strategy to perform assessments for depression in the primary care setting and to improve PHQ-9 scores for patients who are diagnosed with depression. The goal is to improve PHQ-9 scores for patients with depression. The measurable outcome is the percentage of patients with major depression or dysthymia and an initial PHQ-9 score greater than nine whose six-month PHQ-9 score is less than five.
- Sanford Health will continue to provide a directory of services for community members.

**Community Health Needs Assessment
Implementation Strategy for Sanford Hillsboro Medical Center
FY 2017-2020 Action Plan**

Priority 1: Physical health of the community

Projected Impact: Programs and services are available to all members of the community

Goal 1: Provide on-line program Sanford *fit* for children and parents in the community

| Actions/Tactics | Measureable Outcomes | Dedicated Resources | Leadership | Note any community partnerships and collaborations (if applicable) |
|---------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------|-------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------|
| Provide Sanford <i>fit</i> program to the local schools and childcare providers | Sanford <i>fit</i> is available to all students and families in the area through classroom and <i>fit</i> website Will have a kickoff event that introduces the WebMD <i>fit</i> program to the school in late summer. | Sanford <i>fit</i> leadership, classroom teachers, hospital dietitian and managers | Jac McTaggart, Kathy Skager and Kristen Schultz | Hillsboro High School, Central Valley High School, Traill County Public Health, local school nurses and teachers, child care providers |

Goal 2: Exercise programs are available to members of the community

| Actions/Tactics | Measureable Outcomes | Resources | Leadership | Note any community partnerships and collaborations - if applicable |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|-------------------------------|----------------------------------------------------------------------------|
| Work closely with the therapy department and the local wellness center to develop exercise programs (walking clubs, biking clubs, fitness challenges...) for the community | Will have a kick off that will coincide with the Mayville kick off as to work closely with county wide initiatives. We will establish outcomes once we are more into this. We can expand or reduce as we see participation. | Leadership to work with city leaders for utilization of city sidewalks, county roads, Hillsboro Fitness Center, city parks | Kristen Schultz, Kathy Skager | Hillsboro High School, Hillsboro Running Club, Traill County Public Health |

Priority 2: Mental Health Services

Projected Impact: Reduction in the severity of depression

Goal 1: Improve PHQ-9 scores for patients with depression

| Actions/Tactics | Measureable Outcomes | Resources | Leadership | Note any community partnerships and collaborations - if applicable |
|---------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------|-----------------|--------------------------------------------------------------------|
| Develop Sanford My Chart capabilities for depression assessment | Percentage of patients with major depression or dysthymia and an initial PHQ-9 score greater than nine whose six-month PHQ-9 score was less than five. | Mallory Koshiol | Dr. Heidi Twedt | Fist Link |
| Provide education on workflow to all health coaches and panel specialists to standardize workflow | All health coaches in primary care receive education on workflow. | Mallory Koshiol All health coaches | Dr. Heidi Twedt | |

Goal 2: Create awareness of area resources for community members

| Actions/Tactics | Measureable Outcomes | Resources | Leadership | Note any community partnerships and collaborations - if applicable |
|-----------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|-------------------------------------------------------|--------------------------------------------------------------------|
| Distribute directory of available resources to local outreach groups and entities | Need to update and distribute. Will need to work closely with Traill County Health District on distribution. | Sanford Hillsboro, Sanford Health Data Dept., Traill County Health District | Cece Cotton, Mark Duncan, Kathy Skager, Jac McTaggart | Sanford Health Data Department |

2013 Implementation Strategy Impact

2013 Implementation Strategy: Mental Health Services

- Define services currently available
- Define mechanisms to educate service area
- Define education process and secure outside resources
- Develop directory with resources and outsource information
- Distribute directory to various groups, entities and secure email addresses and updates
- Participate in Sanford One Mind as determined by the enterprise three-year plan

2013 Implementation Strategy: Services for the Elderly

- Update directory of available services
- Print/distribute

2013 Implementation Strategy: Day Care

- Determine specific needs through collaboration with county agencies
- Request assistance from outside services to secure daycare services
- Find location for services
- Request community assistance (i.e. keeping daycare profitable)
- Ongoing process - monitoring the needs and services

The 2013 implementation strategies have served a broad base of our community and region. The impact has been positive and the work will continue into the future through new or continued programming and services.

Impact of the Strategy to Address Mental Health Services

Sanford Health completed an inventory of services that are available in the area to address mental health needs. A directory was developed for the purpose of making the information available to the members of the community. The directory was updated in 2015 and will be updated again in 2016.

Revised copies of the mental health resource directory have been made available to public health and community members can obtain a copy at the hospital. Copies will also be distributed to area senior centers during 2016 and at the health fair/education blitz.

Impact of the Strategy to Address Services for the Elderly

Sanford Health completed an inventory of services that are available to meet the needs of seniors in the community and rural areas. A directory was developed for the purpose of making the information available to the members of the community. The directory was updated in 2015, and was added to the mental health services directory.

Revised copies of the mental health and services for the elderly resource directory have been made available to public health and community members can obtain a copy at the hospital. Copies will also be distributed to area senior centers during 2016 and at the health fair/education blitz.

Impact of the Strategy to Address Day Care

Sanford Health worked in collaboration with community leaders to secure daycare services for the community. A director was hired and a local facility was secured and renovated for the daycare services. The provision of day care services in the community fulfilled a need in the community. Late in the fall of 2015 the daycare closed due to the ability to retain and hire new staff. The community is looking at a possible solution of a partnership with the City of Hillsboro and Mayville State to assume ownership of the building and expand the daycare services that Mayville currently offers in Hillsboro.

**Community Feedback
from the 2013
Community Health Needs
Assessment**

Sanford Health is prepared to accept feedback on our 2013 community health needs assessment and has provided on-line comment fields for ease of access on our website. There have been no comments to date aside from a question asked about the service area for this report. A reader wanted to know if a separate report was developed for the Lisbon, North Dakota area. Since there is no hospital in Lisbon, a community health needs assessment was not conducted solely for that community.

APPENDIX

Primary Research

Hillsboro/ Mayville 2016 CHNA Asset Map

| Identified concern | Key stakeholder Survey | Secondary data | Specific areas of concern | Community resources that are available to address the need | Gap? |
|---------------------------|------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Aging population | 4.43 – 3.60 | | <ul style="list-style-type: none"> • Cost of long term care • Availability of memory care • Availability of long term care • Availability of resources for family/friends caring for and making decisions for elders • Availability of activities for seniors • Availability of resources to help the elderly stay safe in their homes | Sanford Home Care – 701-788-4400 Sanford Faith Community Nursing – 701-234-5818 | X |
| Children and Youth | 4.13 – 3.72 | | <ul style="list-style-type: none"> • Availability of activities for children and youth • Availability of quality childcare • Bullying • Availability of quality infant care • Cost of quality child care • Availability of services for at-risk youth • Cost of quality infant care • Cost of activities for children and youth • Cost of services for at-risk youth | NE Human Service Center – 701-795-3000 Teen Line – 1-877-419-7233 (24-hr. counselor line) | X |
| Safety | 3.70 – 3.53 | | <ul style="list-style-type: none"> • Presence of street drugs, and alcohol in the community • Child abuse and neglect • Domestic violence | | X |
| Health Care | 4.09 – 3.61 | | <ul style="list-style-type: none"> • Access to affordable health insurance • Access to affordable health care • Cost of affordable dental insurance coverage • Cost of affordable vision insurance • Access to affordable prescription drugs • Timely access to substance abuse providers • Timely access to mental health providers | Sanford Community Care Program Financial assistance/counselors in business office? | X |
| Physical Health | 4.11 – 3.65 | <ul style="list-style-type: none"> • Reporting poor or fair health – 12% in Steele and 13% in Traill • The obesity rate is 36% in Traill County and 32% in Steele County | <ul style="list-style-type: none"> • Cancer • Chronic Disease (hypertension, high cholesterol, arthritis, diabetes) • Inactivity and lack of exercise (39% have moderate exercise 3 or more times/week, and 17% have vigorous activity 3 or more times/week • Poor nutrition and eating habits (Only 28% have 3 or more vegetables/d, and only 17% have 3 or more fruits/d) • Obesity (65.3% have BMI of | Sanford Cancer Biology Research Center in SF Roger Maris Cancer Center Sanford Medical Home RN Health Coach American Cancer Society Sanford Dietitians | X |

| Identified concern | Key stakeholder Survey | Secondary data | Specific areas of concern | Community resources that are available to address the need | Gap? |
|-----------------------------------------------------------|------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| | | <ul style="list-style-type: none"> The inactivity rate is 27% in Traill County and 30% in Steele County – with a 64% access to exercise opportunities in Traill County and a 27% access in Steel County. | overweight or obese) | <p>Better Choices/Better Health</p> <p>American Heart Association</p> <p>Sanford Clinic – 701-788-4500</p> <p>American Diabetes Association</p> <p>Diabetes Support Group</p> | |
| Mental Health/ Behavioral Health (Substance Abuse) | 3.98 – 3.69 | <ul style="list-style-type: none"> Excessive drinking is at 16% in Trail County and 26% in Steele County 33% of traffic deaths were alcohol impaired in Traill County and 50% were alcohol impaired in Steele County 13% of adults smoke in Traill County and 12% in Steel County | <ul style="list-style-type: none"> Stress (37% report Anxiety/stress) Alcohol use and abuse (29.7% report consuming 3 or more drinks/d and 26.5% have binge level drinking at least 1time/mos.) Underage drinking Depression (31.5% report depression – 55.8% have 1 or more days/mos. when their mental health was not good) Under age drug use and abuse Smoking and tobacco use Drug use and abuse | <p>Sanford Clinic – 701-788-4500</p> <p>NE Human Service Center – 701-795-3000</p> | X |
| Preventive Health | | | <ul style="list-style-type: none"> Flus shots (77% of respondents have had a flu shot and report that 53.8% of children age 6 month or older have had a flu shot) Immunizations (22% of respondents report having immunizations in the past year, and respondents report that 88% of their children are current on their immunizations) 21% have not seen a health care provider in the past year 17% have not seen a dentist in the past year | | |

Hillsboro 2016 Community Health Needs Assessment Prioritization Worksheet

Criteria to Identify Priority Problem

- Cost and/or return on investment
- Availability of solutions
- Impact of problem
- Availability of resources (staff, time, money, equipment) to solve problem
- Urgency of solving problem (H1N1 or air pollution)
- Size of problem (e.g. # of individuals affected)

Criteria to Identify Intervention for Problem

- Expertise to implement solution
- Return on investment
- Effectiveness of solution
- Ease of implementation/maintenance
- Potential negative consequences
- Legal considerations
- Impact on systems or health
- Feasibility of intervention

| Health Indicator/Concern | Round 1 Vote | Round 2 Vote | Round 3 Vote |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------|--------------|--------------|
| Aging <ul style="list-style-type: none"> • Cost of long term care 4.43 • Availability of memory care 3.89 • Availability of LTC 3.81 • Availability of resources for caregivers 3.70 • Availability of activities for seniors 3.61 • Availability of resources to help seniors stay in the homes 3.60 | X | | |
| Children and Youth <ul style="list-style-type: none"> • Availability of activities for children and youth 4.13 • Cost and Availability of quality childcare 4.04 and 3.87 • Bullying 4.04 • Cost and Availability of quality infant care 4.02 and 3.83 • Cost and availability of services for at-risk youth 3.85 and 3.72 • Cost of activities for children and youth 3.80 | | | |
| Safety <ul style="list-style-type: none"> • Seat belt usage • Presence of street drugs and alcohol in the community 3.70 • Child abuse and neglect 3.61 • Domestic violence 3.53 | | | |
| Health care <ul style="list-style-type: none"> • Access to affordable health insurance 4.09 • Access to affordable health care 3.80 • Cost of affordable dental insurance coverage 3.78 • Cost of affordable vision insurance 3.72 • Access to affordable prescription drugs 3.70 • Timely access to substance abuse providers 3.69 • Timely access to mental health providers 3.61 | XX | | |
| Physical Health <ul style="list-style-type: none"> • Cancer 4.11 • Chronic disease 3.98 • Inactivity and lack of exercise 3.81 • Poor nutrition and eating habits 3.70 • Obesity 3.65 • BMI – overweight or obese 65.3% • Only 28.4% of respondents have 3 or more vegetables/day and 17 % have 3 or more fruits/day • Only 39% have 3 or more days each week of moderate activity and 17 % report 3 or more days of vigorous activity each week • 37.7% have smoked at least 100 cigarettes in their life • 22.2% of respondents reported hypertension • 13% reported high cholesterol • 11.1% reported diabetes • Preventive Health – Flus shots and immunization | XXXX #2 priority | | |

| Health Indicator/Concern | Round 1 Vote | Round 2 Vote | Round 3 Vote |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------|--------------|--------------|
| Mental Health <ul style="list-style-type: none"> • Stress 3.98 • Depression 3.81 • 37% of respondents report that they have been told by a doctor that they have anxiety or stress, and 31.5% report being told that they have depression • 55.8% reported 1 or more days in the last month when their mental health was not good. • 29.7% of respondents reported 3 or more drinks /d on average • 26.5% reported 4 or 5 drinks (binge) on the same occasion over the past month • 4.3% reported having a problem with alcohol use; however, 32.7% reported that alcohol use had harmful effects on the respondent or a family member | XXXXX #1 priority | | |

Present: Jill Borth, Steele County Board of Health, DeAnna Kylo, Mayville State University – COP Health, Brittany Ness, Steele County Public Health, Bonnie Overby, Mayville State University, Cassandra Garza, Brenda Stallman, Traill District Health, Jessica Speath, Community of Halstad, Ernie Strube, MayPort EDC, Bill Krivarchka, Eastern ND AHEC, Jac McTaggart, CEO Sanford Hillsboro Medical Center, Jeff McInnes, Sanford Hillsboro Board of Directors, Roger Baier, CEO Sanford Mayville Medical Center

Sanford
Hillsboro/Mayville/Halstad/Trails/Steele
Medical Center

Community Health Needs Assessment
Results from an April 2015 Non-generalizable
Online Survey

August 2015

STUDY DESIGN and METHODOLOGY

The following report includes non-generalizable survey results from an April 2015 on-line survey conducted through a partnership between the Community Health Collaborative and the Center for Social Research (CSR) at North Dakota State University. The CSR developed and maintained links to the on-line survey tool. Members of the Community Health Collaborative invited viewers to access the on-line survey by distributing the survey link via e-mail to various agencies, at times using a snowball approach. **Therefore, it is important to note that the data in this report are not generalizable to the community.** Data collection occurred throughout the month of April 2015 and a total of 54 respondents participated in the on-line survey.

TABLE OF CONTENTS

| | |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| SURVEY RESULTS | 66 |
| General Health and Wellness Concerns about the Community | 66 |
| Figure 1. Level of concern with statements about the community regarding ECONOMICS | |
| Figure 2. Level of concern with statements about the community regarding TRANSPORTATION | |
| Figure 3. Level of concern with statements about the community regarding the ENVIRONMENT | |
| Figure 4. Level of concern with statements about the community regarding CHILDREN AND YOUTH | |
| Figure 5. Level of concern with statements about the community regarding the AGING POPULATION | |
| Figure 6. Level of concern with statements about the community regarding SAFETY | |
| Figure 7. Level of concern with statements about the community regarding HEALTH CARE | |
| Figure 8. Level of concern with statements about the community regarding PHYSICAL AND MENTAL HEALTH | |
| Figure 9. Level of concern with statements about the community regarding SUBSTANCE USE AND ABUSE | |
| General Health | 72 |
| Figure 10. Respondents' rating of their health in general | |
| Figure 11. Respondents' weight status based on the Body Mass Index (BMI) scale | |
| Figure 12. Number of servings of vegetables, fruit, and fruit juice that respondents had yesterday | |
| Figure 13. Number of days in an average week respondents engage in MODERATE and VIGOROUS activity | |
| Mental Health | 75 |
| Figure 14. Percentage of respondents who have been told by a doctor or health professional that they have a mental health issue, by type of mental health issue | |

Figure 15. Number of days in the last month that respondents' mental health was not good

Figure 16. How often, over the past two weeks, respondents have been bothered by mental health issues

Tobacco Use 77

Figure 17. Whether respondents have smoked at least 100 cigarettes in their entire life

Figure 18. How often respondents currently smoke cigarettes and use chewing tobacco or snuff

Figure 19. Location respondents would first go if they wanted help to quit using tobacco

Alcohol Use and Prescription Drug /Non-prescription Drug Abuse 79

Figure 20. Number of days during the past month that respondents had at least one drink of any alcoholic beverage

Figure 21. During the past month on days that respondents drank, average number of drinks per day respondents consumed

Figure 22. Number of times during the past month that respondents consumed at least 4 or 5 alcoholic drinks (4 for females, 5 for males) on the same occasion

Figure 23. Whether respondents have ever had a problem with alcohol use or prescription or non-prescription drug abuse

Figure 24. Of respondents who ever had a problem with alcohol use or prescription or non-prescription drug abuse, whether respondents got the help they needed

Figure 25. Whether alcohol use or prescription or non-prescription drug abuse has had harmful effects on respondents or a family member over the past two years

Preventive Health..... 84

Table 1. Whether or not respondents had preventive screenings in the past year, by type of screening

Table 2. Of respondents who have not had preventive screenings in the past year, reasons why they have not, by type of screening

Figure 26. Whether respondents have any of the following chronic diseases

Figure 27. Length of time since respondents last visited a doctor or health care provider for a routine physical exam and length of time since they last visited a dentist or dental clinic for any reason

Figure 28. Where respondents get most of their health information

Figure 29. Best way for respondents to access technology for health information

Demographic Information..... 89

Figure 30. Age of respondents

Figure 31. Highest level of education of respondents

Figure 32. Gender of respondents

Figure 33. Race and ethnicity of respondents

Figure 34. Annual household income of respondents

Figure 35. Employment status of respondents

Figure 36. Length of time respondents have lived in their community

Figure 37. Whether respondents own or rent their home

Figure 38. Whether respondents have health insurance (private, public, or governmental) and oral health or dental care insurance coverage

Figure 39. Whether respondents have one person who they think of as their personal doctor or health care provider

Figure 40. Facilities that respondents go to most often when sick and take their children when they are sick

Figure 41. Number of children younger than 18 and number of adults age 65 or older living in respondents' household

Figure 42. Whether all children in home are current on their immunizations and all children age 6 months or older get a flu shot or flu mist each year

Table 3. Zip code of respondents

SURVEY RESULTS

General Health and Wellness Concerns about the Community

Using a 1 to 5 scale, with 1 being “not at all” and 5 being “a great deal,” respondents were asked to rate their level of concern with various statements regarding ECONOMICS, TRANSPORTATION, the ENVIRONMENT, CHILDREN AND YOUTH, the AGING POPULATION, SAFETY, HEALTH CARE, PHYSICAL AND MENTAL HEALTH, and SUBSTANCE USE AND ABUSE.

Figure 1. Level of concern with statements about the community regarding ECONOMICS




Figure 2. Level of concern with statements about the community regarding TRANSPORTATION




Figure 3. Level of concern with statements about the community regarding the ENVIRONMENT




Figure 4. Level of concern with statements about the community regarding CHILDREN AND YOUTH




Figure 5. Level of concern with statements about the community regarding the AGING POPULATION




Figure 6. Level of concern with statements about the community regarding SAFETY




Figure 7. Level of concern with statements about the community regarding HEALTH CARE




Figure 8. Level of concern with statements about the community regarding PHYSICAL AND MENTAL HEALTH






Figure 9. Level of concern with statements about the community regarding SUBSTANCE USE AND ABUSE




General Health

Figure 10. Respondents' rating of their health in general




N=53 Figure 11. Respondents' weight status based on the Body Mass Index (BMI)* scale



N=49


*For information about the BMI, visit the Centers for Disease Control and Prevention, *About BMI for Adults*, http://www.cdc.gov/healthyweight/assessing/bmi/adult_bmi/.

Figure 12. Number of servings of vegetables, fruit, and fruit juice that respondents had yesterday



*Percentages may not total 100.0 due to rounding.


Figure 13. Number of days in an average week respondents engage in MODERATE and VIGOROUS activity



*Percentages do not total 100.0 due to rounding.


Mental Health

Figure 14. Percentage of respondents who have been told by a doctor or health professional that they have a mental health issue, by type of mental health issue




N=54 *Percentages do not total 100.0 due to multiple responses.

Figure 15. Number of days in the last month that respondents' mental health was not good



N=43


Figure 16. How often, over the past two weeks, respondents have been bothered by mental health issues



*Percentages may not total 100.0 due to rounding.

Tobacco Use

Figure 17. Whether respondents have smoked at least 100 cigarettes in their entire life



N=53

Figure 18. How often respondents currently smoke cigarettes and use chewing tobacco or snuff






Figure 19. Location respondents would first go if they wanted help to quit using tobacco



N=53 *Other response is "medical center".


Alcohol Use and Prescription Drug/Non-prescription Drug Abuse

Figure 20. Number of days during the past month that respondents had at least one drink of any alcoholic beverage



N=50


Figure 21. During the past month on days that respondents drank, average number of drinks per day respondents consumed



N=37

*Percentages do not total 100.0 due to rounding.

Figure 22. Number of times during the past month that respondents consumed at least 4 or 5 alcoholic drinks (4 for females, 5 for males) on the same occasion



N=53

*Percentages do not total 100.0 due to rounding.

Figure 23. Whether respondents ever had a problem with alcohol use or prescription or non-prescription drug abuse




Figure 24. Of respondents who ever had a problem with alcohol use or prescription or non-prescription drug abuse, whether respondents got the help they needed





Figure 25. Whether alcohol use or prescription or non-prescription drug abuse has had harmful effects on respondents or a family member over the past two years



Preventive Health

Table 1. Whether or not respondents had preventive screenings in the past year, by type of screening


| Type of screening | Percent of respondents | | |
|------------------------------------------|------------------------|-------|-------|
| | Yes | No | Total |
| GENERAL SCREENINGS | | | |
| Blood pressure screening (N=52) | 86.5 | 13.5 | 100.0 |
| Blood sugar screening (N=51) | 64.7 | 35.3 | 100.0 |
| Bone density test (N=50) | 6.0 | 94.0 | 100.0 |
| Cardiovascular screening (N=50) | 24.0 | 76.0 | 100.0 |
| Cholesterol screening (N=52) | 67.3 | 32.7 | 100.0 |
| Dental screening and X-rays (N=52) | 82.7 | 17.3 | 100.0 |
| Flu shot (N=52) | 76.9 | 23.1 | 100.0 |
| Glaucoma test (N=52) | 50.0 | 50.0 | 100.0 |
| Hearing screening (N=49) | 6.1 | 93.9 | 100.0 |
| Immunizations (N=50) | 22.0 | 78.0 | 100.0 |
| Pelvic exam (N=45 Females) | 60.0 | 40.0 | 100.0 |
| STD (N=50) | 12.0 | 88.0 | 100.0 |
| Vascular screening (N=51) | 11.8 | 88.2 | 100.0 |
| CANCER SCREENINGS | | | |
| Breast cancer screening (N=45 Females) | 75.6 | 24.4 | 100.0 |
| Cervical cancer screening (N=45 Females) | 60.0 | 40.0 | 100.0 |
| Colorectal cancer screening (N=50) | 18.0 | 82.0 | 100.0 |
| Prostate cancer screening (N=5 Males) | 0.0 | 100.0 | 100.0 |
| Skin cancer screening (N=49) | 14.3 | 85.7 | 100.0 |

Table 2. Of respondents who have not had preventive screenings in the past year, reasons why they have not, by type of screening

| Type of screening | Percentage of respondents* | | | | | | |
|------------------------------------------|----------------------------|-------------------------|------|-------------------|-----------------|-----------------------|--------------|
| | Not necessary | Doctor hasn't suggested | Cost | Fear of procedure | Fear of results | Unable to access care | Other reason |
| GENERAL SCREENINGS | | | | | | | |
| Blood pressure screening (N=7) | 71.4 | 28.6 | 0.0 | 0.0 | 0.0 | 14.3 | 0.0 |
| Blood sugar screening (N=18) | 61.1 | 33.3 | 11.1 | 0.0 | 0.0 | 0.0 | 0.0 |
| Bone density test (N=47) | 55.3 | 31.9 | 8.5 | 0.0 | 0.0 | 0.0 | 0.0 |
| Cardiovascular screening (N=38) | 52.6 | 36.8 | 7.9 | 0.0 | 0.0 | 2.6 | 0.0 |
| Cholesterol screening (N=17) | 64.7 | 41.2 | 5.9 | 0.0 | 0.0 | 0.0 | 0.0 |
| Dental screening and (X-rays (N=9) | 22.2 | 22.2 | 66.7 | 22.2 | 0.0 | 11.1 | 11.1 |
| Flu shot (N=12) | 41.7 | 8.3 | 8.3 | 0.0 | 0.0 | 0.0 | 50.0 |
| Glaucoma test (N=26) | 65.4 | 23.1 | 7.7 | 0.0 | 0.0 | 3.8 | 0.0 |
| Hearing screening (N=46) | 60.9 | 28.3 | 6.5 | 0.0 | 0.0 | 2.2 | 2.2 |
| Immunizations (N=39) | 74.4 | 17.9 | 2.6 | 0.0 | 0.0 | 0.0 | 2.6 |
| Pelvic exam (N=18 Females) | 66.7 | 27.8 | 0.0 | 0.0 | 0.0 | 0.0 | 11.1 |
| STD (N=44) | 79.5 | 15.9 | 2.3 | 2.3 | 0.0 | 0.0 | 0.0 |
| Vascular screening (N=45) | 57.8 | 28.9 | 8.9 | 0.0 | 2.2 | 4.4 | 0.0 |
| CANCER SCREENINGS | | | | | | | |
| Breast cancer screening (N=11 Females) | 63.6 | 36.4 | 9.1 | 0.0 | 0.0 | 0.0 | 0.0 |
| Cervical cancer screening (N=18 Females) | 61.1 | 16.7 | 5.6 | 0.0 | 0.0 | 0.0 | 5.6 |
| Colorectal cancer screening (N=41) | 53.7 | 22.0 | 4.9 | 12.2 | 0.0 | 2.4 | 4.9 |
| Prostate cancer screening (N=5 Males) | 40.0 | 20.0 | 20.0 | 20.0 | 0.0 | 20.0 | 20.0 |
| Skin cancer screening (N=42) | 47.6 | 38.1 | 7.1 | 2.4 | 0.0 | 2.4 | 4.8 |

*Percentages may not total 100.0 due to multiple responses.


Figure 26. Whether respondents have any of the following chronic diseases



N=54


*Percentages do not total 100.0 due to multiple responses.

Figure 27. Length of time since respondents last visited a doctor or health care provider for a routine physical exam and length of time since they last visited a dentist or dental clinic for any reason




*Percentages may not total 100.0 due to rounding.

Figure 28. Where respondents get most of their health information



N=54 *Percentages do not total 100.0 due to multiple responses. **Other responses include “by working in health care”, “co-workers”, and “the internet”.


Figure 29. Best way for respondents to access technology for health information



N=54 *Percentages do not total 100.0 due to multiple responses. ** Other response is “provider’s ofc.”


Demographic Information

Figure 30. Age of respondents



N=52


Figure 31. Highest level of education of respondents



N=52


*Percentages do not total 100.0 due to rounding

Figure 32. Gender of respondents



N=52


Figure 33. Race and ethnicity of respondents




N=54

*Percentages do not total 100.0 due to multiple responses

Figure 34. Annual household income of respondents




N=52 Figure 35. Employment status of respondents



N=52


Figure 36. Length of time respondents have lived in their community



N=52

*Percentages do not total 100.0 due to rounding.

Figure 37. Whether respondents own or rent their home



N=52

Figure 38. Whether respondents have health insurance (private, public, or governmental) and oral health or dental care insurance coverage






Figure 39. Whether respondents have one person who they think of as their personal doctor or health care provider



N=52


Fig. 40. Facilities that respondents go to most often when sick & take their children when they are sick



*Percentages may not total 100.0 due to rounding.


**Of respondents who have children younger than 18 years living in their household.

Figure 41. Number of children younger than 18 and number of adults age 65 or older living in respondents' household



*Percentages may not total 100.0 due to rounding.

Figure 42. Whether all children in home are current on their immunizations and all children age 6 months or older get a flu shot or flu mist each year*



*Of respondents who have children younger than 18 years living in their household

Table 3. Zip code of respondents

| Zip Code | Number of respondents |
|-----------------|------------------------------|
| 58257 | 13 |
| 58230 | 9 |
| 58274 | 6 |
| 58045 | 5 |
| 58240 | 4 |
| 56548 | 3 |
| 58016 | 3 |
| 58218 | 3 |
| 55060 | 1 |
| 58035 | 1 |
| 58038 | 1 |
| 58046 | 1 |
| 58103 | 1 |

N=51

Secondary Research

Trail County Community Health Profile

April 13, 2015

Stephen Pickard, MD

North Dakota Department of Health

TABLE OF CONTENTS

POPULATION DATA

| | |
|--------------------------------------------------------------------|---|
| Chart 1: Population by Age Group | 1 |
| Chart 2: Pyramid Graph of Number of Residents by Census Year | 1 |
| Chart 3: Female Population and Percentage Female by Age | 1 |
| Chart 4: Decennial Population Change | 1 |
| Chart 5: Race | 2 |
| Chart 6: Household Populations | 2 |
| Chart 7: Marital Status of Persons Age 15 and Older | 2 |
| Chart 8: Educational Attainment Among Persons 25+ | 2 |
| Chart 9: Persons with Disability | 3 |
| Chart 10: Income and Poverty Status by Age Group | 3 |
| Chart 11: Family Poverty and Childhood and Elderly Poverty | 3 |
| Chart 12: Age of Housing | 3 |

VITAL STATISTICS DATA

| | |
|------------------------------------------------------------------------|---|
| Birth and Deaths Definitions | 4 |
| Chart 13: Births | 5 |
| Chart 14: Child Deaths | 5 |
| Chart 15: Deaths and Age Adjusted Death Rate by Cause | 5 |
| Chart 16: Leading Causes of Death by Age Group for Traill County | 6 |
| Chart 17: Leading Causes of Death by Age Group for North Dakota | 6 |

ADULT BEHAVIORAL RISK FACTORS

| | |
|--------------------------------------------------|----|
| Chart 18 –21: | |
| • Adult Behavioral Risk Factors Definition | 7 |
| • Alcohol | 8 |
| • Arthritis | 8 |
| • Asthma | 8 |
| • Body Weight | 8 |
| • Cancer | 8 |
| • Cardiovascular | 9 |
| • Cholesterol | 9 |
| • Chronic Lung Disease | 9 |
| • Colorectal Cancer | 9 |
| • Diabetes | 9 |
| • Fruits and Vegetables | 9 |
| • General Health | 10 |
| • Health Care Access | 10 |
| • Hypertension | 10 |
| • Immunization | 10 |
| • Injury | 10 |
| • Oral Health | 11 |
| • Physical Activity | 11 |
| • Tobacco | 11 |
| • Women’s Health | 11 |

CRIME

| | |
|-------------------------------|----|
| Chart 21: Traill County | 12 |
| Chart 22: North Dakota | 12 |

CHILD RISK


| | |
|----------------------------------------------------------|----|
| Chart 23: Child Indicators—Education | 13 |
| Chart 24: Child Indicators—Economic Health | 13 |
| Chart 25: Child Indicators—Families and Child Care | 13 |
| Chart 26: Child Indicators—Juvenile Justice | 13 |

POPULATION DATA

1

| Population by Age Group, 2009-2013 ACS Estimates | | | | |
|--------------------------------------------------|---------------|---------------|----------------|---------------|
| Age Group | Traill County | | North Dakota | |
| | Number | Percent | Number | Percent |
| 0-9 | 963 | 11.9% | 91,871 | 12.7% |
| 10-19 | 1,075 | 13.2% | 93,318 | 12.9% |
| 20-29 | 973 | 12.0% | 124,424 | 17.2% |
| 30-39 | 852 | 10.5% | 91,148 | 12.6% |
| 40-49 | 1,001 | 12.3% | 79,573 | 11.0% |
| 50-59 | 1,197 | 14.7% | 101,998 | 14.1% |
| 60-69 | 909 | 11.2% | 69,446 | 9.6% |
| 70-79 | 576 | 7.1% | 41,233 | 5.7% |
| 80+ | 579 | 7.1% | 31,829 | 4.4% |
| Total | 8,125 | 100.0% | 724,840 | 100.0% |
| 0-17 | 1,786 | 22.0% | 161,317 | 22.3% |
| 65+ | 1,511 | 18.6% | 102,722 | 14.2% |

The Demographic Section of this report comes from the US Census Bureau (www.census.gov). Most tables are derived either from the Census estimates for 2013 or from the Community Population Survey aggregated over a several year period. The table header describes the specific years from which the data is derived. The table showing percent population change uses census data from 2000 also. Tables present number of persons and percentages which in almost all circumstances represent the category specific percentage of all persons referenced by the table (e.g., percentage of persons age 15 and older who are married). Age specific poverty rates represent the percentage of each age group which is in poverty (e.g., percentage of children under five years in poverty).



3

| Female Population and Percentage Female by Age, 2009-2013 ACS Estimates | | | | |
|-------------------------------------------------------------------------|---------------|--------------|----------------|--------------|
| Age Group | Traill County | | North Dakota | |
| | Number | Percent | Number | Percent |
| 0-9 | 453 | 47.0% | 45,342 | 49.4% |
| 10-19 | 534 | 49.7% | 44,988 | 48.2% |
| 20-29 | 449 | 46.1% | 55,615 | 44.7% |
| 30-39 | 425 | 49.9% | 41,092 | 45.1% |
| 40-49 | 478 | 47.8% | 39,675 | 49.9% |
| 50-59 | 557 | 46.5% | 50,656 | 49.7% |
| 60-69 | 477 | 52.5% | 34,361 | 49.5% |
| 70-79 | 301 | 52.3% | 21,254 | 51.5% |
| 80+ | 381 | 65.8% | 20,546 | 64.5% |
| Total | 4,055 | 49.9% | 353,530 | 48.8% |
| 0-17 | 865 | 48.4% | 78,484 | 48.7% |
| 65+ | 882 | 58.4% | 50,201 | 48.9% |

4

| Decennial Population Change, 1990 to 2000, 2000 to 2010 | | | | |
|---------------------------------------------------------|---------------|--------------------|--------------|--------------------|
| Census | Traill County | 10 Year Change (%) | North Dakota | 10 Year Change (%) |
| 1990 | 8,752 | | 638,800 | |
| 2000 | 8,477 | -3.1% | 642,200 | 0.5% |
| 2010 | 8,121 | -4.2% | 672,591 | 4.7% |

POPULATION DATA

9

| Persons with Disability, 2009-2013 ACS Estimates | | | | |
|--------------------------------------------------|---------------|---------|--------------|---------|
| Group | Traill County | | North Dakota | |
| | Number | Percent | Number | Percent |
| Total | 7,984 | 100.0% | 688,158 | 100.0% |
| Any Disability | 855 | 10.7% | 72,762 | 10.6% |
| No Disability | 7,129 | 89.3% | 615,396 | 89.4% |
| Self Care Disability | 156 | 2.0% | 11,274 | 1.6% |
| 0-17 with any disability | 66 | 3.7% | 4,677 | 2.9% |
| 18-64 with any disability | 402 | 8.3% | 35,931 | 7.8% |
| 65+ with any disability | 387 | 28.1% | 32,154 | 31.3% |

10

| Income and Poverty Status by Age Group, 2009-2013 ACS Estimates | | | | |
|-----------------------------------------------------------------|---------------|---------|--------------|---------|
| | Traill County | | North Dakota | |
| | Number | Percent | Number | Percent |
| Median Household Income | \$48,767 | | \$54,920 | |
| Per Capita Income | \$27,096 | | \$30,436 | |
| Below Poverty Level | 828 | 10.7% | 80,644 | 11.9% |
| Under 5 years | 21 | 17.4% | 7,714 | 16.9% |
| 5 to 11 years | 125 | 17.4% | 7,944 | 13.2% |
| 12 to 17 years | 48 | 8.3% | 5,776 | 11.8% |
| 18 to 64 years | 427 | 9.2% | 49,568 | 11.6% |
| 65 to 74 years | 52 | 7.6% | 3,448 | 7.0% |
| 75 years and over | 110 | 15.8% | 6,194 | 13.7% |
| Total Known Children in Poverty | 194 | 10.9% | 21,434 | 13.3% |
| Total Known Age 65+ in Poverty | 162 | 10.7% | 9,642 | 9.4% |

11

| Family Poverty, 2009-2013 ACS Estimates | | | | |
|--------------------------------------------------------------|---------------|----------|--------------|----------|
| | Traill County | | North Dakota | |
| | Number | Percent* | Number | Percent* |
| Total Families | 1,964 | 100.0% | 176,378 | 100.0% |
| Families in Poverty | 132 | 6.7% | 13,052 | 7.4% |
| Families with Own Children | 842 | 42.9% | 80,964 | 45.9% |
| Families with Own Children in Poverty | 103 | 5.2% | 10,121 | 5.7% |
| Families with Own Children and Female Parent Only | 144 | 7.3% | 16,716 | 9.5% |
| Families with Own Children and Female Parent Only in Poverty | 80 | 4.1% | 6,452 | 3.7% |

12

| Age of Housing, 2009-2013 ACS Estimates | | | | |
|-----------------------------------------|---------------|---------|--------------|---------|
| | Traill County | | North Dakota | |
| | Number | Percent | Number | Percent |
| Housing units: Total | 3,778 | 100.0% | 329,970 | 100.0% |
| 1980 and Later | 815 | 21.6% | 128,111 | 38.8% |
| 1970 to 1979 | 748 | 19.8% | 66,396 | 20.1% |
| Prior to 1970 | 2,215 | 58.6% | 135,463 | 41.1% |

Vital Statistics Data

BIRTHS AND DEATHS DEFINITIONS

Vital Statistics Data comes from the birth and death records collected by the State of North Dakota aggregated over a five year period. All births and deaths represent the county of residence not the county of occurrence. The number of events is blocked if fewer than five. Formulas for calculating rates and ratios are as follows:

Birth Rate = Resident live births divided by the total resident population x 1000.

Pregnancies = Live births + Fetal deaths + Induced termination of pregnancy.

Pregnancy Rate = Total pregnancies divided by the total resident population x 1000.

Fertility Rate = Resident live births divided by female population (age 15-44) x 1000.

Teenage Birth Rate = Teenage births (age <20) divided by female teen population x 1000.

Teenage Pregnancy Rate =
Teenage pregnancies (age <20) divided by female teen population x 1000.

Out of Wedlock Live Birth Ratio =
Resident OOW live births divided by total resident live births x 1000.

Out of Wedlock Pregnancy Ratio =
Resident OOW pregnancies divided by total pregnancies x 1000.

Low Weight Ratio =
Low weight births (birth weight < 2500 grams) divided by total resident live births x 1000.

Infant Death Ratio = Number of infant deaths divided by the total resident live births x 1000.

Childhood & Adolescent Deaths = Deaths to individuals 1 - 19 years of age.

Childhood and Adolescent Death Rate =
Number of resident deaths (age 1 - 19) divided by population (age 1 - 19) x 100,000.

Crude Death Rate = Death events divided by population x 100,000.

Age-Adjusted Death Rate = Death events with age specific adjustments x 100,000 population.

Vital Statistics Data

BIRTHS AND DEATHS

13

| | Traill County | | North Dakota | |
|----------------------------------|---------------|---------------|--------------|---------------|
| | Number | Rate or Ratio | Number | Rate or Ratio |
| Births, 2009-2013 | | | | |
| Live Births and Rate | 474 | 12 | 47,959 | 14 |
| Pregnancies and Rate | 502 | 12 | 52,505 | 15 |
| Fertility Rate | | 70 | | 72 |
| Teen Births and Rate | 23 | 11 | 2,118 | 12 |
| Teen Pregnancies and Rate | 29 | 14 | 3,725 | 21 |
| Out of Wedlock Births and Ratio | 131 | 276 | 15,686 | 327 |
| Out of Wedlock Preg and Ratio | 156 | 311 | 19,436 | 370 |
| Low Birth Weight Birth and Ratio | 8 | 17 | 3,078 | 64 |

14

| | Traill County | | North Dakota | |
|--------------------------------------|---------------|---------------|--------------|---------------|
| | Number | Rate or Ratio | Number | Rate or Ratio |
| Child Deaths, 2009-2013 | | | | |
| Infant Deaths and Ratio | * | NA | 286 | 6 |
| Child and Adolescent Deaths and Rate | * | NA | 270 | 32 |
| Total Deaths and Crude Rate | 448 | 1,103 | 29,616 | 866 |

15

| | Traill County | | North Dakota | |
|---------------------------------------------------------------|--------------------|--------------------|--------------------|--------------------|
| | Number (Adj. Rate) | Number (Adj. Rate) | Number (Adj. Rate) | Number (Adj. Rate) |
| Deaths and Age Adjusted Death Rate by Cause, 2009-2013 | | | | |
| All Causes | 448 (661) | | 29,581 (702) | |
| Heart Disease | 133 (169) | | 6,762 (154) | |
| Cancer | 89 (154) | | 6,315 (156) | |
| Stroke | 36 (44) | | 1,664 (37) | |
| Alzheimers Disease | 31 (37) | | 2,189 (45) | |
| COPD | 24 (36) | | 1,707 (41) | |
| Unintentional Injury | 17 (34) | | 1,625 (44) | |
| Diabetes Mellitus | 9 (13) | | 1,022 (24) | |
| Pneumonia and Influenza | 9 (13) | | 682 (15) | |
| Cirrhosis | * | | 394 (11) | |
| Suicide | * | | 551 (16) | |

Vital Statistics Data

BIRTHS AND DEATHS

16


| Leading Causes of Death by Age Group for Traill County, 2009-2013 | | | |
|-------------------------------------------------------------------|----------------------|--------------|-----------------------------------|
| Age | 1 | 2 | 3 |
| 0-4 | Congenital Anomaly | | |
| 5-14 | | | |
| 15-24 | Unintentional Injury | | |
| 25-34 | Unintentional Injury | | |
| 35-44 | Cancer | Suicide | Unintentional Injury Cirrhosis |
| 45-54 | Cancer | Heart | Unintentional Injury |
| 55-64 | Cancer 6 | Heart | COPD |
| 65-74 | Cancer 10 | Heart 7 | Stroke 5 |
| 75-84 | Cancer 19 | Heart 12 | COPD 11 |
| 85+ | Cancer 32 | Heart 31 | Alzheimer's Dz 21 |
| | Heart 81 | Stroke 24 | |

17

| Leading Causes of Death by Age Group for North Dakota, 2006-2010 | | | |
|------------------------------------------------------------------|-----------------------------|------------------------|-----------------------------|
| Age | 1 | 2 | 3 |
| 0-4 | Congenital Anomaly 69 | Prematurity 44 | SIDS 40 |
| 5-14 | Unintentional Injury 26 | Cancer 10 | Congenital Anomaly 6 |
| 15-24 | Unintentional Injury 184 | Suicide 109 | Cancer 20 |
| 25-34 | Unintentional Injury 166 | Suicide 91 | Heart 32 |
| 35-44 | Unintentional Injury 173 | Heart 94 | Cancer 88 |
| 45-54 | Cancer 493 | Heart 335 | Unintentional Injury 194 |
| 55-64 | Cancer 1001 | Heart 579 | Unintentional Injury 137 |
| 65-74 | Cancer 1562 | Heart 843 | COPD 313 |
| 75-84 | Cancer 1992 | Heart 1797 | COPD 626 |
| 85+ | Heart 3421 | Alzheimer's Dz 1391 | Cancer 1352 |

ADULT BEHAVIORAL RISK FACTORS DEFINITION

The following three pages represent data received from the Adult Behavioral Risk Factor Surveillance Survey. The Adult Behavioral Risk Factor data are derived from aggregated data (the number of years specified is in the table) continuously collected by telephone survey from persons 18 years and older. All data is self-reported data. Numbers given are point estimate percentages followed by 95% confidence intervals. Statistical significance can be determined by comparing confidence intervals between two geographic areas. To be statistically significant, confidence intervals may not overlap. For example the confidence intervals 9.3 (8.3-10.2) and 10.8 (10.0-11.6) overlap (see picture below) so the difference between the two numbers is not statistically significant. That means that substantial uncertainty remains whether the apparent difference is due to chance alone (due to sampling variation) rather than representing a true difference in the prevalence of the condition in the two populations. The less they overlap, the more likely it is that the point estimates represent truly different prevalences in the two populations.



8...9...10...11...12...

ADULT BEHAVIORAL RISK FACTORS, 2011-2013

18

| ALCOHOL | | | | |
|--------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------|---------------------------------------------|---------------------------|
| | | Trail and Steele 2011-2013 | Trail, Steele, Sargent, Ransom 2011-2013 | North Dakota 2011-2013 |
| Binge Drinking | Respondents who reported binge drinking (5 drinks for men, 4 drinks for women) one or more times in the past 30 days. | 20.3 (12.4-28.3) | 24.3 (18.8-29.9) | 24.2 (23.2-25.1) |
| Heavy Drinking | Respondents who reported heavy drinking (more than 2 drinks per day for men, more than 1 drink per day for women) during the past 30 days | NA | 7.5 (4.3-10.7) | 6.8 (6.2-7.4) |
| Drunk Driving | Respondents who reported driving when they had too much to drink one or more times during the past 30 days | NA | NA | 1.9 (1.5-2.4) |
| ARTHRITIS | | | | |
| Doctor Diagnosed Arthritis | Respondents who reported ever have been told by a doctor or other health professional that they had some form of arthritis. | 24.2 (18.5-29.9) | 28.8 (24.4-33.3) | 25.1 (24.3-25.9) |
| Activity Limitation Due to Arthritis | Respondents who reported being limited in any usual activities because of arthritis or joint symptoms. | NA | 9.5 (6.6-12.4) | 8.3 (7.8-8.8) |
| ASTHMA | | | | |
| Ever Asthma | Respondents who reported ever having been told by a doctor, nurse or other health professional that they had asthma. | 10.0 (5.3-14.7) | 10.6 (7.1-14.2) | 11.5 (10.7-12.2) |
| Current Asthma | Respondents who reported ever having been told by a doctor, nurse or other health professional that they had asthma and who still have asthma. | NA | 8.2 (4.9-11.6) | 8.1 (7.5-8.7) |
| BODY WEIGHT | | | | |
| Overweight But Not Obese | Respondents with a body mass index greater than or equal to 25 but less than 30 (overweight) | 39.0 (30.2-47.9) | 37.8 (32.0-43.7) | 36.7 (35.7-37.7) |
| Obese | Respondents with a body mass index greater than or equal to 30 (obese) | 37.2 (28.1-46.2) | 33.8 (28.1-39.5) | 29.4 (28.5-30.4) |
| Overweight or Obese | Respondents with a body mass index greater than or equal to 25 (overweight or obese) | 76.2 (68.2-84.2) | 71.6 (65.9-77.3) | 66.1 (65.1-67.2) |
| CANCER | | | | |
| Ever Cancer | Respondents who reported ever having been told by a doctor, nurse or other health professional that they had cancer (excluding skin cancer). | 9.0 (5.3-12.7) | 7.5 (5.2-9.9) | 6.4 (6.0-6.8) |

ADULT BEHAVIORAL RISK FACTORS, 2011-2013

20

| | GENERAL HEALTH | Trails and Steele 2011-2013 | Trails, Steele, Sargent, Ransom 2011-2013 | North Dakota 2011-2013 |
|----------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------|----------------------------------------------|---------------------------|
| Fair or Poor Health | Respondents who reported that their general health was fair or poor | 14.0 (8.8-19.2) | 11.8 (8.6-14.9) | 14.0 (13.3-14.7) |
| Poor physical Health | Respondents who reported they had 8 or more days in the last 30 when their physical health was not good | 10.5 (5.6-15.4) | 9.1 (5.8-12.4) | 11.6 (11.0-12.3) |
| Poor Mental Health | Respondents who reported they had 8 or more days in the last 30 when their mental health was not good | 11.2 (4.9-17.4) | 10.3 (6.2-14.4) | 10.8 (10.2-11.5) |
| Activity Limitation Due to Poor Health | Respondents who reported they had 8 or more days in the last 30 when poor physical or mental health kept them from doing their usual activities. | NA | NA | 7.1 (6.6-7.6) |
| Any Activity Limitation | Respondents who reported being limited in any way due to physical, mental or emotional problem. | 17.3 (11.7-22.9) | 15.2 (11.7-18.7) | 17.9 (17.2-18.7) |
| HEALTH CARE ACCESS | | | | |
| Health Insurance | Respondents who reported not having any form or health care coverage | NA | 9.0 (5.5-12.4) | 12.4 (11.6-13.1) |
| Access Limited by Cost | Respondents who reported needing to see a doctor during the past 12 months but could not due to cost. | NA | NA | 8.1 (7.4-8.7) |
| No Personal Provider | Respondents who reported that they did not have one person they consider to be their personal doctor or health care provider. | 20.7 (12.8-28.6) | 20.4 (15.5-25.4) | 26.0 (25.0-26.9) |
| HYPERTENSION | | | | |
| High Blood Pressure | Respondents who reported ever having been told by a doctor, nurse or other health professional that they had high blood pressure. | 24.0 (16.7-31.2) | 28.9 (23.5-34.3) | 29.5 (28.6-30.6) |
| IMMUNIZATION | | | | |
| Influenza Vaccine | Respondents age 65 and older who reported that they did not have a flu shot in the past year | 24.2 (14.5-33.9) | 32.5 (24.2-40.9) | 40.3 (38.6-42.0) |
| Pneumococcal Vaccine | Respondents age 65 or older who reported never having had a pneumonia shot. | NA | 28.2 (19.6-36.8) | 29.4 (27.8-31.0) |
| INJURY | | | | |
| Fall | Respondents 45 years and older who reported that they had fallen in the past 12 months | NA | 28.2 (19.0-37.5) | 27.8 (25.9-29.6) |
| Seat Belt | Respondents who reported not always wearing their seatbelt | 45.7 (37.1-54.3) | 47.8 (41.9-53.7) | 32.6 (31.6-33.6) |

ADULT BEHAVIORAL RISK FACTORS, 2011-2013

21

| | ORAL HEALTH | Traill and Steele 2011-2013 | Traill, Steele, Sargent, Ransom 2011-2013 | North Dakota 2011-2013 |
|-----------------------------|--------------------------------------------------------------------------------------------|--------------------------------|----------------------------------------------|---------------------------|
| Dental Visit | Respondents who reported that they have not had a dental visit in the past year | NA | NA | 32.8 (31.0-34.7) |
| Tooth Loss | Respondents who reported they ever had a permanent tooth extracted. | NA | NA | 43.2 (41.4-45.0) |
| PHYSICAL ACTIVITY | | | | |
| Recommend Physical Activity | Respondents who reported that they did not get the recommended amount of physical activity | NA | 51.7 (44.6-58.8) | 53.8 (52.6-55.1) |
| Inactive | Respondents reporting little or no physical activity | 24.8 (15.8-33.9) | 29.7 (23.4-36.0) | 30.4 (29.3-31.6) |
| TOBACCO | | | | |
| Current Smoking | Respondents who reported that they smoked every day or some days | 17.4 (10.4-24.5) | 18.0 (13.4-22.5) | 21.6 (20.6-22.5) |
| WOMEN'S HEALTH | | | | |
| Pap Smear | Women 18 and older who reported that they have not had a pap smear in the past three years | NA | NA | 23.7 (20.9-26.5) |
| Mammogram Age 40+ | Women 40 and older who reported that they have not had a mammogram in the past two years | NA | NA | 26.4 (24.0-28.9) |

Steele County Community Health Profile

April 13, 2015



Stephen Pickard, MD
North Dakota Department of Health

TABLE OF CONTENTS

POPULATION DATA

| | |
|-------------------------------------------------------------------|---|
| Chart 1: Population by Age Group..... | 1 |
| Chart 2: Pyramid Graph of Number of Residents by Census Year..... | 1 |
| Chart 3: Female Population and Percentage Female by Age..... | 1 |
| Chart 4: Decennial Population Change..... | 1 |
| Chart 5: Race..... | 2 |
| Chart 6: Household Populations..... | 2 |
| Chart 7: Marital Status of Persons Age 15 and Older..... | 2 |
| Chart 8: Educational Attainment Among Persons 25+..... | 2 |
| Chart 9: Persons with Disability..... | 3 |
| Chart 10: Income and Poverty Status by Age Group..... | 3 |
| Chart 11: Family Poverty and Childhood and Elderly Poverty..... | 3 |
| Chart 12: Age of Housing..... | 3 |

VITAL STATISTICS DATA

| | |
|-----------------------------------------------------------------------|---|
| Birth and Deaths Definitions..... | 4 |
| Chart 13: Births..... | 5 |
| Chart 14: Child Deaths..... | 5 |
| Chart 15: Deaths and Age Adjusted Death Rate by Cause..... | 5 |
| Chart 16: Leading Causes of Death by Age Group for Steele County..... | 6 |
| Chart 17: Leading Causes of Death by Age Group for North Dakota..... | 6 |

ADULT BEHAVIORAL RISK FACTORS

| | |
|-------------------------------------------------|----|
| Chart 18 –21: | |
| • Adult Behavioral Risk Factors Definition..... | 7 |
| • Alcohol..... | 8 |
| • Arthritis..... | 8 |
| • Asthma..... | 8 |
| • Body Weight..... | 8 |
| • Cancer..... | 8 |
| • Cardiovascular..... | 9 |
| • Cholesterol..... | 9 |
| • Chronic Lung Disease..... | 9 |
| • Colorectal Cancer..... | 9 |
| • Diabetes..... | 9 |
| • Fruits and Vegetables..... | 9 |
| • General Health..... | 10 |
| • Health Care Access..... | 10 |
| • Hypertension..... | 10 |
| • Immunization..... | 10 |
| • Injury..... | 10 |
| • Oral Health..... | 11 |
| • Physical Activity..... | 11 |
| • Tobacco..... | 11 |
| • Women’s Health..... | 11 |

CRIME

| | |
|------------------------------|----|
| Chart 21: Steele County..... | 12 |
| Chart 22: North Dakota..... | 12 |

CHILD RISK

| | |
|---------------------------------------------------------|----|
| Chart 23: Child Indicators—Education..... | 13 |
| Chart 24: Child Indicators—Economic Health..... | 13 |
| Chart 25: Child Indicators—Families and Child Care..... | 13 |
| Chart 26: Child Indicators—Juvenile Justice..... | 13 |


POPULATION DATA

1

| Age Group | Steele County | | North Dakota | |
|--------------|---------------|---------------|----------------|---------------|
| | Number | Percent | Number | Percent |
| 0-9 | 203 | 10.3% | 91,871 | 12.7% |
| 10-19 | 254 | 12.9% | 93,318 | 12.9% |
| 20-29 | 166 | 8.4% | 124,424 | 17.2% |
| 30-39 | 208 | 10.5% | 91,148 | 12.6% |
| 40-49 | 261 | 13.2% | 79,573 | 11.0% |
| 50-59 | 375 | 19.0% | 101,998 | 14.1% |
| 60-69 | 223 | 11.3% | 69,446 | 9.6% |
| 70-79 | 174 | 8.8% | 41,233 | 5.7% |
| 80+ | 112 | 5.7% | 31,829 | 4.4% |
| Total | 1,976 | 100.0% | 724,840 | 100.0% |
| 0-17 | 424 | 21.5% | 161,317 | 22.3% |
| 65+ | 376 | 19.0% | 102,722 | 14.2% |

The Demographic Section of this report comes from the US Census Bureau (www.census.gov). Most tables are derived either from the Census estimates for 2013 or from the Community Population Survey aggregated over a several year period. The table header describes the specific years from which the data is derived. The table showing percent population change uses census data from 2000 also. Tables present number of persons and percentages which in almost all circumstances represent the category specific percentage of all persons referenced by the table (e.g., percentage of persons age 15 and older who are married). Age specific poverty rates represent the percentage of each age group which is in poverty (e.g., percentage of children under five years in poverty).

2



3

| Age Group | Steele County | | North Dakota | |
|--------------|---------------|--------------|----------------|--------------|
| | Number | Percent | Number | Percent |
| 0-9 | 98 | 48.3% | 45,342 | 49.4% |
| 10-19 | 125 | 49.2% | 44,988 | 48.2% |
| 20-29 | 71 | 42.8% | 55,615 | 44.7% |
| 30-39 | 90 | 43.3% | 41,092 | 45.1% |
| 40-49 | 139 | 53.3% | 39,675 | 49.9% |
| 50-59 | 166 | 44.3% | 50,656 | 49.7% |
| 60-69 | 113 | 50.7% | 34,361 | 49.5% |
| 70-79 | 97 | 55.7% | 21,254 | 51.5% |
| 80+ | 51 | 45.5% | 20,546 | 64.5% |
| Total | 950 | 48.1% | 353,530 | 48.8% |
| 0-17 | 112 | 26.4% | 78,484 | 48.7% |
| 65+ | 186 | 49.5% | 50,201 | 48.9% |

4

| Census | Steele County | | North Dakota | |
|--------|--------------------|--------------------|--------------------|--------------------|
| | 10 Year Change (%) |
| 1990 | 2,420 | | 638,800 | |
| 2000 | 2,258 | -6.7% | 642,200 | 0.5% |
| 2010 | 1,975 | -12.5% | 672,591 | 4.7% |

POPULATION DATA

5

| Race, 2009-2013 ACS Estimates | | | | |
|-------------------------------|---------------|------------|--------------|------------|
| Race | Steele County | | North Dakota | |
| | Number | Percentage | Number | Percentage |
| Total | 1,976 | 100.0% | 723,393 | 100.0% |
| White | 1,944 | 98.4% | 643,478 | 89.0% |
| Black | 0 | 0.0% | 10,827 | 1.5% |
| Am. Indian | 0 | 0.0% | 40,214 | 5.6% |
| Asian | 0 | 0.0% | 9,096 | 1.3% |
| Pac. Islander | 0 | 0.0% | 371 | 0.1% |
| Other | 32 | 1.6% | 4,620 | 0.6% |
| Multirace | 0 | 0.0% | 29,574 | 4.1% |

6

| Household Populations, 2011-2013 ACS Three Year Estimates | | | | |
|-----------------------------------------------------------|---------------|---------|--------------|---------|
| | Steele County | | North Dakota | |
| | Number | Percent | Number | Percent |
| Total | 1,976 | 100.0% | 703,203 | 100.0% |
| In Family Households | 1,618 | 81.9% | 530,615 | 75.5% |
| In Non-Family Households | 358 | 18.1% | 146,330 | 20.8% |
| Total In Households | 1,976 | 100.0% | 676,945 | 96.3% |
| Institutionalized* | 0 | 0.0% | 9,675 | 1.4% |
| Non-institutionalized* | 0 | 0.0% | 16,583 | 2.4% |
| Total in Group Quarters | 0 | 0.0% | 26,258 | 3.7% |

7

| Marital Status of Persons Age 15 and Older, 2009-2013 ACS Estimates | | | | |
|---------------------------------------------------------------------|---------------|---------|--------------|---------|
| Marital Status | Steele County | | North Dakota | |
| | Number | Percent | Number | Percent |
| Total Age 15+ | 1,631 | 100.0% | 561,346 | 100.0% |
| Never Married | 287 | 17.6% | 177,385 | 31.6% |
| Now Married | 1,073 | 65.8% | 291,900 | 52.0% |
| Separated | 7 | 0.4% | 5,052 | 0.9% |
| Widowed | 101 | 6.2% | 33,681 | 6.0% |
| Divorced | 161 | 9.9% | 53,328 | 9.5% |

8

| Educational Attainment Among Persons 25+, 2009-2013 ACS Estimates | | | | |
|-------------------------------------------------------------------|---------------|---------|--------------|---------|
| Education | Steele County | | North Dakota | |
| | Number | Percent | Number | Percent |
| Total | 1,421 | 100.0% | 457,771 | 100.0% |
| Less than 9th Grade | 53 | 3.7% | 19,226 | 4.2% |
| Some High School | 63 | 4.4% | 21,057 | 4.6% |
| High school or GRE | 441 | 31.0% | 125,429 | 27.4% |
| Some College / Asso. Degree | 580 | 40.8% | 168,002 | 36.7% |
| Bachelor's degree | 217 | 15.3% | 89,723 | 19.6% |
| Post Graduate Degree | 70 | 4.9% | 34,791 | 7.6% |

POPULATION DATA

9

| Persons with Disability, 2009-2013 ACS Estimates | | | | |
|--------------------------------------------------|---------------|---------|--------------|---------|
| Group | Steele County | | North Dakota | |
| | Number | Percent | Number | Percent |
| Total | 1,976 | 100.0% | 688,158 | 100.0% |
| Any Disability | 204 | 10.3% | 72,762 | 10.6% |
| No Disability | 1,772 | 89.7% | 615,396 | 89.4% |
| Self Care Disability | 22 | 1.2% | 11,274 | 1.6% |
| 0-17 with any disability | 6 | 1.4% | 4,677 | 2.9% |
| 18-64 with any disability | 97 | 8.2% | 35,931 | 7.8% |
| 65+ with any disability | 101 | 26.9% | 32,154 | 31.3% |

10

| Income and Poverty Status by Age Group, 2009-2013 ACS Estimates | | | | |
|-----------------------------------------------------------------|---------------|---------|--------------|---------|
| | Steele County | | North Dakota | |
| | Number | Percent | Number | Percent |
| Median Household Income | \$55,735 | | \$54,920 | |
| Per Capita Income | \$31,455 | | \$30,436 | |
| Below Poverty Level | 121 | 6.2% | 80,644 | 11.9% |
| Under 5 years | 9 | 9.8% | 7,714 | 16.9% |
| 5 to 11 years | 7 | 4.9% | 7,944 | 13.2% |
| 12 to 17 years | 4 | 2.5% | 5,776 | 11.8% |
| 18 to 64 years | 47 | 4.0% | 49,568 | 11.6% |
| 65 to 74 years | 20 | 11.3% | 3,448 | 7.0% |
| 75 years and over | 34 | 17.1% | 6,194 | 13.7% |
| Total Known Children in Poverty | 20 | 4.7% | 21,434 | 13.3% |
| Total Known Age 65+ in Poverty | 54 | 14.4% | 9,642 | 9.4% |

11

| Family Poverty and Childhood and Elderly Poverty, 2009-2013 ACS Estimates | | | | |
|---------------------------------------------------------------------------|---------------|----------|--------------|----------|
| | Steele County | | North Dakota | |
| | Number | Percent* | Number | Percent* |
| Total Families | 595 | 100.0% | 176,378 | 100.0% |
| Families in Poverty | 24 | 4.0% | 13,052 | 7.4% |
| Families with Own Children | 210 | 35.3% | 80,964 | 45.9% |
| Families with Own Children in Poverty | 19 | 3.2% | 10,121 | 5.7% |
| Families with Own Children and Female Parent Only | 38 | 6.4% | 16,716 | 9.5% |
| Families with Own Children and Female Parent Only in Poverty | 15 | 2.5% | 6,452 | 3.7% |

* Percent family poverty is percent of total families

12

| Age of Housing ,2009-2013 ACS Estimates | | | | |
|-----------------------------------------|---------------|---------|--------------|---------|
| | Steele County | | North Dakota | |
| | Number | Percent | Number | Percent |
| Housing units: Total | 1,152 | 100.0% | 329,970 | 100.0% |
| 1980 and Later | 243 | 21.1% | 128,111 | 38.8% |
| 1970 to 1979 | 179 | 15.5% | 66,396 | 20.1% |
| Prior to 1970 | 730 | 63.4% | 135,463 | 41.1% |

Vital Statistics Data

BIRTHS AND DEATHS DEFINITIONS

Vital Statistics Data comes from the birth and death records collected by the State of North Dakota aggregated over a five year period. All births and deaths represent the county of residence not the county of occurrence. The number of events is blocked if fewer than five. Formulas for calculating rates and ratios are as follows:

Birth Rate = Resident live births divided by the total resident population x 1000.

Pregnancies = Live births + Fetal deaths + Induced termination of pregnancy.

Pregnancy Rate = Total pregnancies divided by the total resident population x 1000.

Fertility Rate = Resident live births divided by female population (age 15-44) x 1000.

Teenage Birth Rate = Teenage births (age <20) divided by female teen population x 1000.

Teenage Pregnancy Rate =
Teenage pregnancies (age <20) divided by female teen population x 1000.

Out of Wedlock Live Birth Ratio =
Resident OOW live births divided by total resident live births x 1000.

Out of Wedlock Pregnancy Ratio =
Resident OOW pregnancies divided by total pregnancies x 1000.

Low Weight Ratio =
Low weight births (birth weight < 2500 grams) divided by total resident live births x 1000.

Infant Death Ratio = Number of infant deaths divided by the total resident live births x 1000.

Childhood & Adolescent Deaths = Deaths to individuals 1 - 19 years of age.

Childhood and Adolescent Death Rate =
Number of resident deaths (age 1 - 19) divided by population (age 1 - 19) x 100,000.

Crude Death Rate = Death events divided by population x 100,000.

Age-Adjusted Death Rate = Death events with age specific adjustments x 100,000 population.

Vital Statistics Data

BIRTHS AND DEATHS

13

| | Steele County | | North Dakota | |
|----------------------------------|---------------|---------------|--------------|---------------|
| | Number | Rate or Ratio | Number | Rate or Ratio |
| Births, 2009-2013 | | | | |
| Live Births and Rate | 92 | 9 | 47,959 | 14 |
| Pregnancies and Rate | 96 | 10 | 52,505 | 15 |
| Fertility Rate | | 67 | | 72 |
| Teen Births and Rate | 6 | 13 | 2,118 | 12 |
| Teen Pregnancies and Rate | 7 | 15 | 3,725 | 21 |
| Out of Wedlock Births and Ratio | 24 | 261 | 15,686 | 327 |
| Out of Wedlock Preg and Ratio | 26 | 271 | 19436 | 370 |
| Low Birth Weight Birth and Ratio | * | NA | 3,078 | 64 |

14

| | Steele County | | North Dakota | |
|--------------------------------------|---------------|---------------|--------------|---------------|
| | Number | Rate or Ratio | Number | Rate or Ratio |
| Child Deaths, 2009-2013 | | | | |
| Infant Deaths and Ratio | 0 | 0 | 286 | 6 |
| Child and Adolescent Deaths and Rate | * | NA | 270 | 32 |
| Total Deaths and Crude Rate | 84 | 850 | 29,616 | 866 |

15

| | Steele County | | North Dakota | |
|---------------------------------------------------------------|--------------------|--------------------|--------------------|--------------------|
| | Number (Adj. Rate) | Number (Adj. Rate) | Number (Adj. Rate) | Number (Adj. Rate) |
| Deaths and Age Adjusted Death Rate by Cause, 2009-2013 | | | | |
| All Causes | 84 (587) | | 29,581 (702) | |
| Heart Disease | 20 (129) | | 6,762 (154) | |
| Cancer | 20 (137) | | 6,315 (156) | |
| Stroke | 6 (36) | | 1,664 (37) | |
| Alzheimers Disease | 6 (33) | | 2,189 (45) | |
| COPD | 5 (31) | | 1,707 (41) | |
| Unintentional Injury | 8 (90) | | 1,625 (44) | |
| Diabetes Mellitus | * | | 1,022 (24) | |
| Pneumonia and Influenza | 0 | | 682 (15) | |
| Cirrhosis | 0 | | 394 (11) | |
| Suicide | * | | 551 (16) | |

Vital Statistics Data

BIRTHS AND DEATHS

16

| Leading Causes of Death by Age Group for Steele County, 2009-2013 | | | |
|-------------------------------------------------------------------|----------------------|---------------------|----------------------|
| Age | 1 | 2 | 3 |
| 0-4 | | | |
| 5-14 | Cancer | | |
| 15-24 | Unintentional Injury | | |
| 25-34 | Unintentional Injury | | |
| 35-44 | Suicide | | |
| 45-54 | Heart | Cancer | Unintentional Injury |
| 55-64 | Cancer | | |
| 65-74 | Cancer 5 | Heart | |
| 75-84 | Cancer 7 Heart 7 | COPD Stroke | Unintentional Injury |
| 85+ | Heart 7 | Alzheimer's Dz 5 | Cancer Stroke |

17

| Leading Causes of Death by Age Group for North Dakota, 2006-2010 | | | |
|------------------------------------------------------------------|-----------------------------|------------------------|-----------------------------|
| Age | 1 | 2 | 3 |
| 0-4 | Congenital Anomaly 69 | Prematurity 44 | SIDS 40 |
| 5-14 | Unintentional Injury 26 | Cancer 10 | Congenital Anomaly 6 |
| 15-24 | Unintentional Injury 184 | Suicide 109 | Cancer 20 |
| 25-34 | Unintentional Injury 166 | Suicide 91 | Heart 32 |
| 35-44 | Unintentional Injury 173 | Heart 94 | Cancer 88 |
| 45-54 | Cancer 493 | Heart 335 | Unintentional Injury 194 |
| 55-64 | Cancer 1001 | Heart 579 | Unintentional Injury 137 |
| 65-74 | Cancer 1562 | Heart 843 | COPD 313 |
| 75-84 | Cancer 1992 | Heart 1797 | COPD 626 |
| 85+ | Heart 3421 | Alzheimer's Dz 1391 | Cancer 1352 |

ADULT BEHAVIORAL RISK FACTORS DEFINITION

The following three pages represent data received from the Adult Behavioral Risk Factor Surveillance Survey. The Adult Behavioral Risk Factor data are derived from aggregated data (the number of years specified is in the table) continuously collected by telephone survey from persons 18 years and older. All data is self-reported data. Numbers given are point estimate percentages followed by 95% confidence intervals. Statistical significance can be determined by comparing confidence intervals between two geographic areas. To be statistically significant, confidence intervals may not overlap. For example the confidence intervals 9.3 (8.3-10.2) and 10.8 (10.0-11.6) overlap (see picture below) so the difference between the two numbers is not statistically significant. That means that substantial uncertainty remains whether the apparent difference is due to chance alone (due to sampling variation) rather than representing a true difference in the prevalence of the condition in the two populations. The less they overlap, the more likely it is that the point estimates represent truly different prevalences in the two populations.

8.....9.....10.....11.....12.....

ADULT BEHAVIORAL RISK FACTORS, 2011-2013

19

| | CARDIOVASCULAR | Trail and Steele 2011-2013 | Trail, Steele, Sargent, Ransom 2011-2013 | North Dakota 2011-2013 |
|-------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------|-------------------------------------------------------------|---------------------------------------|
| Heart Attack | Respondents who reported ever having been told by a doctor, nurse or other health care professional that they had a heart attack. | NA | 4.0 (2.5- 5.5) | 4.3 (3.9-4.6) |
| Angina | Respondents who reported ever having been told by a doctor, nurse or other health care professional that they had angina. | NA | 3.3 (1.9- 4.7) | 4.1 (3.7-4.4) |
| Stroke | Respondents who reported ever having been told by a doctor, nurse or other health care professional that they had a stroke. | NA | NA | 2.3 (2.0-2.5) |
| Cardiovascular Disease | Respondents who reported ever having been told by a doctor, nurse or other health care professional that they had any of the following: heart attack, angina or stroke. | 6.2 (3.3- 9.1) | 6.5 (4.5- 8.4) | 7.6 (7.1-8.0) |
| CHOLESTEROL | | | | |
| Never Cholesterol Test | Respondents who reported never having a cholesterol test | NA | 19.0 (12.6-25.4) | 22.3 (21.1-23.4) |
| No Cholesterol Test in Past 5 Years | Respondents who reported never having a cholesterol test in the past five years | NA | 24.6 (18.0-31.1) | 26.7 (25.5-27.9) |
| High Cholesterol | Respondents who reported that they had ever been told by a doctor, nurse or other health professional that they had high cholesterol. | 40.3 (30.2-50.4) | 40.3 (33.8-46.9) | 36.6 (35.4-37.8) |
| CHRONIC LUNG DISEASE | | | | |
| COPD | Respondents who have ever been told by a doctor, nurse or other health professional ever told you that they have COPD (chronic obstructive pulmonary disease), emphysema, or chronic bronchitis? | NA | 5.7 (3.6- 7.9) | 4.6 (4.2-5.0) |
| COLORECTAL CANCER | | | | |
| Fecal Occult Blood | Respondents age 50 and older who reported not having a fecal occult blood test in the past two years. | 90.0 (81.6-98.4) | 81.5 (73.2-89.9) | 86.2 (84.8-87.6) |
| Never Sigmoidoscopy | Respondents age 50 and older who reported never having had a sigmoidoscopy or colonoscopy | NA | 45.2 (33.4-56.9) | 38.0 (35.9-40.2) |
| Up to date for Colorectal Screening | Respondents age 50 and older who are up to date according to recommended screening guidelines for colorectal screening | NA | 53.8 (42.1-65.4) | 59.1 (56.9-61.3) |
| DIABETES | | | | |
| Diabetes Diagnosis | Respondents who reported ever having been told by a doctor that they had diabetes. | 10.5 (6.1-14.8) | 9.9 (7.0-12.8) | 8.5 (8.0-9.0) |
| FRUITS AND VEGETABLES | | | | |
| Five Fruits and Vegetables | Respondents who reported that they do not usually eat 5 fruits and vegetables per day | 84.4 (76.2-92.7) | 87.5 (82.8-92.2) | 85.9 (85.0-86.7) |

ADULT BEHAVIORAL RISK FACTORS, 2011-2013

20

| | GENERAL HEALTH | Traill and Steele 2011-2013 | Traill, Steele, Sargent, Ransom 2011-2013 | North Dakota 2011-2013 |
|----------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------|-------------------------------------------------------|------------------------------|
| Fair or Poor Health | Respondents who reported that their general health was fair or poor | 14.0 (8.8-19.2) | 11.8 (8.6-14.9) | 14.0 (13.3-14.7) |
| Poor physical Health | Respondents who reported they had 8 or more days in the last 30 when their physical health was not good | 10.5 (5.6-15.4) | 9.1 (5.8-12.4) | 11.6 (11.0-12.3) |
| Poor Mental Health | Respondents who reported they had 8 or more days in the last 30 when their mental health was not good | 11.2 (4.9-17.4) | 10.3 (6.2-14.4) | 10.8 (10.2-11.5) |
| Activity Limitation Due to Poor Health | Respondents who reported they had 8 or more days in the last 30 when poor physical or mental health kept them from doing their usual activities. | NA | NA | 7.1 (6.6-7.6) |
| Any Activity Limitation | Respondents who reported being limited in any way due to physical, mental or emotional problem. | 17.3 (11.7-22.9) | 15.2 (11.7-18.7) | 17.9 (17.2-18.7) |
| HEALTH CARE ACCESS | | | | |
| Health Insurance | Respondents who reported not having any form or health care coverage | NA | 9.0 (5.5-12.4) | 12.4 (11.6-13.1) |
| Access Limited by Cost | Respondents who reported needing to see a doctor during the past 12 months but could not due to cost. | NA | NA | 8.1 (7.4-8.7) |
| No Personal Provider | Respondents who reported that they did not have one person they consider to be their personal doctor or health care provider. | 20.7 (12.8-28.6) | 20.4 (15.5-25.4) | 26.0 (25.0-26.9) |
| HYPERTENSION | | | | |
| High Blood Pressure | Respondents who reported ever having been told by a doctor, nurse or other health professional that they had high blood pressure. | 24.0 (16.7-31.2) | 28.9 (23.5-34.3) | 29.5 (28.6-30.6) |
| IMMUNIZATION | | | | |
| Influenza Vaccine | Respondents age 65 and older who reported that they did not have a flu shot in the past year | 24.2 (14.5-33.9) | 32.5 (24.2-40.9) | 40.3 (38.6-42.0) |
| Pneumococcal Vaccine | Respondents age 65 or older who reported never having had a pneumonia shot. | NA | 28.2 (19.6-36.8) | 29.4 (27.8-31.0) |
| INJURY | | | | |
| Fall | Respondents 45 years and older who reported that they had fallen in the past 12 months | NA | 28.2 (19.0-37.5) | 27.8 (25.9-29.6) |
| Seat Belt | Respondents who reported not always wearing their seatbelt | 45.7 (37.1-54.3) | 47.8 (41.9-53.7) | 32.6 (31.6-33.6) |

CRIME

Steele County

21

| | 2009 | 2010 | 2011 | 2012 | 2013 | 4 year | 4-Year Rate |
|-----------------------|------|----------|----------|----------|----------|-----------|--------------|
| Murder | NA | 0 | 0 | 0 | 0 | 0 | 0.0 |
| Rape | | 0 | 0 | 1 | 0 | 1 | 12.5 |
| Robbery | | 0 | 0 | 0 | 0 | 0 | 0.0 |
| Aggrav. Assault | | 1 | 0 | 0 | 2 | 3 | 37.5 |
| Violent crime | | 1 | 0 | 1 | 2 | 4 | 50.0 |
| Burglary | | 0 | 3 | 5 | 1 | 9 | 112.5 |
| Larceny | | 1 | 2 | 1 | 1 | 5 | 62.5 |
| Motor vehicle theft | | 0 | 1 | 1 | 1 | 3 | 37.5 |
| Property crime | | 1 | 6 | 7 | 3 | 17 | 212.5 |
| Total | | 2 | 6 | 8 | 5 | 21 | 262.5 |

North Dakota

22

| | 2009 | 2010 | 2011 | 2012 | 2013 | 5 year | 5-Year Rate |
|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| Murder | 15 | 11 | 15 | 20 | 14 | 75 | 2.2 |
| Rape | 206 | 222 | 207 | 243 | 237 | 1,115 | 32.6 |
| Robbery | 102 | 85 | 91 | 117 | 151 | 546 | 16.0 |
| Aggrav. Assault | 795 | 847 | 1,040 | 1,071 | 1,156 | 4,909 | 143.6 |
| Violent crime | 1,118 | 1,165 | 1,353 | 1,451 | 1,558 | 6,645 | 194.3 |
| Burglary | 2,180 | 1,826 | 2,227 | 2,200 | 2,656 | 11,089 | 324.3 |
| Larceny | 8,699 | 8,673 | 9,344 | 10,184 | 10,243 | 47,143 | 1378.6 |
| Motor vehicle theft | 854 | 825 | 763 | 854 | 1,228 | 4,524 | 132.3 |
| Property crime | 11,733 | 11,324 | 12,334 | 13,238 | 14,127 | 62,756 | 1835.2 |
| Total | 12,851 | 12,489 | 13,687 | 14,689 | 15,685 | 69,401 | 2029.5 |

Definitions of Key Indicators

**County Health
Rankings & Roadmaps**

Building a Culture of Health, County by County

A Robert Wood Johnson Foundation program

A collaboration between the Robert Wood Johnson Foundation and the University of Wisconsin Population Health Institute. This Excel file contains the ranks and scores for each county in your state and the underlying data details for the measures used in calculating the 2015 *County Health Rankings*. In addition, the file contains additional measures that are reported on the *County Health Rankings* web site for your state.

For additional information about how the *County Health Rankings* are calculated, please visit www.countyhealthrankings.org

Contents:

Outcomes & Factors Rankings

Outcomes & Factors Sub Rankings

Ranked Measures Data (including measure values, confidence intervals* and z-scores**)

Additional Measures Data (including measure values and confidence intervals*)

Ranked Measure Sources and Years

Additional Measure Sources and Years

* 95% confidence intervals are provided where applicable and available.

** Z-scores are "adjusted" z-scores (e.g., multiplied by -1 if a positively framed measure, set to zero for missing and unreliable values for ranked counties, and truncated at -3 or +3 if county population is less than 20,000).

| Measure | Data Elements | Description |
|-------------------------------|------------------------------------------|---------------------------------------------------------------------------|
| Geographic identifiers | FIPS | Federal Information Processing Standard |
| | State | |
| | County | |
| Premature death | # Deaths | Number of deaths under age 75 |
| | Years of Potential Life Lost Rate | Age-adjusted YPLL rate per 100,000 |
| | 95% CI - Low | 95% confidence interval reported by National Center for Health Statistics |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Poor or fair health | Sample Size | Number of respondents |
| | % Fair/Poor | Percent of adults that report fair or poor health |
| | 95% CI - Low | 95% confidence interval reported by BRFSS |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |

| Measure | Data Elements | Description |
|-----------------------------------------|----------------------------------|-----------------------------------------------------------------------------------|
| Poor physical health days | Sample Size | Number of respondents |
| | Physically Unhealthy Days | Average number of reported physically unhealthy days per month |
| | 95% CI - Low | 95% confidence interval reported by BRFSS |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Poor mental health days | Sample Size | Number of respondents |
| | Mentally Unhealthy Days | Average number of reported mentally unhealthy days per month |
| | 95% CI - Low | 95% confidence interval reported by BRFSS |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Low birthweight | Unreliable | Value reported but considered unreliable since based on counts of twenty or less. |
| | # Low Birthweight Births | Number of low birthweight births |
| | # Live births | Number of live births |
| | % LBW | Percentage of births with low birth weight (<2500g) |
| | 95% CI - Low | 95% confidence interval reported by National Center for Health Statistics |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Adult smoking | Sample Size | Number of respondents |
| | % Smokers | Percentage of adults that reported currently smoking |
| | 95% CI - Low | 95% confidence interval reported by BRFSS |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Adult obesity | % Obese | Percentage of adults that report BMI >= 30 |
| | 95% CI - Low | 95% confidence interval reported by BRFSS |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Food environment index | Food Environment Index | Indicator of access to healthy foods - 0 is worst, 10 is best |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Physical inactivity | % Physically Inactive | Percentage of adults that report no leisure-time physical activity |
| | 95% CI - Low | 95% confidence interval |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Access to exercise opportunities | # With Access | Number of people with access to exercise opportunities |
| | % With Access | Percentage of the population with access to places for physical activity |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Excessive drinking | Sample Size | Number of respondents |
| | % Excessive Drinking | Percentage of adults that report excessive drinking |
| | 95% CI - Low | 95% confidence interval reported by BRFSS |

| Measure | Data Elements | Description |
|----------------------------------------|-----------------------------------|--------------------------------------------------------------------------------|
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Alcohol-impaired driving deaths | # Alcohol-Impaired Driving Deaths | Number of alcohol-impaired motor vehicle deaths |
| | # Driving Deaths | Number of motor vehicle deaths |
| | % Alcohol-Impaired | Percentage of driving deaths with alcohol involvement |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Sexually transmitted infections | # Chlamydia Cases | Number of chlamydia cases |
| | Chlamydia Rate | Chlamydia cases / Population * 100,000 |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Teen births | Teen Births | Teen birth count, ages 15-19 |
| | Teen Population | Female population, ages 15-19 |
| | Teen Birth Rate | Teen births / females ages 15-19 * 1,000 |
| | 95% CI - Low | 95% confidence interval reported by National Center for Health Statistics |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Uninsured | # Uninsured | Number of people under age 65 without insurance |
| | % Uninsured | Percentage of people under age 65 without insurance |
| | 95% CI - Low | 95% confidence interval reported by SAHIE |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Primary care physicians | # Primary Care Physicians | Number of primary care physicians (PCP) in patient care |
| | PCP Rate | (Number of PCP/population)*100,000 |
| | PCP Ratio | Population to Primary Care Physicians ratio |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Dentists | # Dentists | Number of dentists |
| | Dentist Rate | (Number of dentists/population)*100,000 |
| | Dentist Ratio | Population to Dentists ratio |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Mental health providers | # Mental Health Providers | Number of mental health providers (MHP) |
| | MHP Rate | (Number of MHP/population)*100,000 |
| | MHP Ratio | Population to Mental Health Providers ratio |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Preventable hospital stays | # Medicare Enrollees | Number of Medicare enrollees |
| | Preventable Hosp. Rate | Discharges for Ambulatory Care Sensitive Conditions/Medicare Enrollees * 1,000 |
| | 95% CI - Low | 95% confidence interval reported by Dartmouth Institute |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Diabetic monitoring | # Diabetics | Number of diabetic Medicare enrollees |
| | % Receiving HbA1c | Percentage of diabetic Medicare enrollees receiving HbA1c |

| Measure | Data Elements | Description |
|---------------------------------------------|-----------------------------------|------------------------------------------------------------------------------------------|
| | | test |
| | 95% CI - Low | 95% confidence interval reported by Dartmouth Institute |
| | 95% CI - High | |
| | Z-Score | |
| Mammography screening | # Medicare Enrollees | Number of female Medicare enrollees age 67-69 |
| | % Mammography | Percentage of female Medicare enrollees having at least 1 mammogram in 2 yrs (age 67-69) |
| | 95% CI - Low | 95% confidence interval reported by Dartmouth Institute |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| High school graduation | Cohort Size | Number of students expected to graduate |
| | Graduation Rate | Graduation rate |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Some college | # Some College | Adults age 25-44 with some post-secondary education |
| | Population | Adults age 25-44 |
| | % Some College | Percentage of adults age 25-44 with some post-secondary education |
| | 95% CI - Low | 95% confidence interval |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Unemployment | # Unemployed | Number of people ages 16+ unemployed and looking for work |
| | Labor Force | Size of the labor force |
| | % Unemployed | Percentage of population ages 16+ unemployed and looking for work |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Children in poverty | # Children in Poverty | Number of children (under age 18) living in poverty |
| | % Children in Poverty | Percentage of children (under age 18) living in poverty |
| | 95% CI - Low | 95% confidence interval reported by SAIPE |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Income inequality | 80th Percentile Income | 80th percentile of median household income |
| | 20th Percentile Income | 20th percentile of median household income |
| | Income Ratio | Ratio of household income at the 80th percentile to income at the 20th percentile |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Children in single-parent households | # Single-Parent Households | Number of children that live in single-parent households |
| | # Households | Number of children in households |
| | % Single-Parent Households | Percentage of children that live in single-parent households |
| | 95% CI - Low | 95% confidence interval |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Social associations | # Associations | Number of associations |
| | Association Rate | Associations / Population * 10,000 |

| Measure | Data Elements | Description |
|-------------------------------------------|--------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------|
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Violent crime | # Violent Crimes | Number of violent crimes |
| | Violent Crime Rate | Violent crimes/population * 100,000 |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Injury deaths | # Injury Deaths | Number of injury deaths |
| | Injury Death Rate | Injury mortality rate per 100,000 |
| | 95% CI - Low | 95% confidence interval as reported by the National Center for Health Statistics |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Air pollution - particulate matter | Average Daily PM2.5 | Average daily amount of fine particulate matter in micrograms per cubic meter |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Drinking water violations | Pop. In Viol | Average annual population affected by a water violation |
| | % Pop in Viol | Population affected by a water violation/Total population with public water |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Severe housing problems | # Households with Severe Problems | Number of households with at least 1 of 4 housing problems: overcrowding, high housing costs, or lack of kitchen or plumbing facilities |
| | % Severe Housing Problems | Percentage of households with at least 1 of 4 housing problems: overcrowding, high housing costs, or lack of kitchen or plumbing facilities |
| | 95% CI - Low | 95% confidence interval |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Driving alone to work | # Drive Alone | Number of people who drive alone to work |
| | # Workers | Number of workers in labor force |
| | % Drive Alone | Percentage of workers who drive alone to work |
| | 95% CI - Low | 95% confidence interval |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |
| Long commute - driving alone | # Workers who Drive Alone | Number of workers who commute in their car, truck or van alone |
| | % Long Commute - Drives Alone | Among workers who commute in their car alone, the percentage that commute more than 30 minutes |
| | 95% CI - Low | 95% confidence interval |
| | 95% CI - High | |
| | Z-Score | (Measure - Average of state counties)/(Standard Deviation) |

Trail County

| | Trail County | Trend(Click for info) | Error Margin | Top U.S. Performers* | North Dakota | Rank (of 47) |
|----------------------------------|--------------|-------------------------------------------------------------------------------------|--------------|----------------------|--------------|--------------|
| Health Outcomes | | | | | | 1 |
| Length of Life | | | | | | 1 |
| Premature death | 4,400 |  | 3,403-5,598 | 5,200 | 6,388 | |
| Quality of Life | | | | | | 7 |
| Poor or fair health | 13% | | 9-18% | 10% | 12% | |
| Poor physical health days | 2.7 | | 1.8-3.5 | 2.5 | 2.7 | |
| Poor mental health days | 2.2 | | 1.2-3.2 | 2.3 | 2.4 | |
| Low birthweight | 4.2% | | 2.7-5.7% | 5.9% | 6.5% | |
| Health Factors | | | | | | 7 |
| Health Behaviors | | | | | | 13 |
| Adult smoking | 13% | | 9-18% | 14% | 18% | |
| Adult obesity | 36% |  | 30-42% | 25% | 30% | |
| Food environment index | 8.9 | | | 8.4 | 8.5 | |
| Physical inactivity | 27% |  | 22-32% | 20% | 25% | |
| Access to exercise opportunities | 64% | | | 92% | 68% | |
| Excessive drinking | 16% | | 12-21% | 10% | 22% | |
| Alcohol-impaired driving deaths | 33% | | | 14% | 46% | |
| Sexually transmitted infections | 161 |  | | 138 | 416 | |
| Teen births | 15 | | 10-21 | 20 | 28 | |

| | Trail County | Trend(Click for info) | Error Margin | Top U.S. Performers* | North Dakota | Rank (of 47) |
|--------------------------------------|--------------|-------------------------------------------------------------------------------------|--------------|----------------------|--------------|--------------|
| Clinical Care | | | | | | 11 |
| Uninsured | 11% |  | 10-13% | 11% | 12% | |
| Primary care physicians | 4,036:1 | | | 1,045:1 | 1,279:1 | |
| Dentists | 4,123:1 | | | 1,377:1 | 1,710:1 | |
| Mental health providers | | | | 386:1 | 638:1 | |
| Preventable hospital stays | 55 |  | 43-66 | 41 | 56 | |
| Diabetic monitoring | 92% |  | 74-100% | 90% | 86% | |
| Mammography screening | 74.3% |  | 57.8-90.8% | 70.7% | 68.0% | |
| Social & Economic Factors | | | | | | 3 |
| High school graduation | | | | | 85% | |
| Some college | 76.4% | | 67.6-85.1% | 71.0% | 74.4% | |
| Unemployment | 3.1% |  | | 4.0% | 2.9% | |
| Children in poverty | 9% |  | 6-12% | 13% | 12% | |
| Income inequality | 4.3 | | 3.7-5.0 | 3.7 | 4.4 | |
| Children in single-parent households | 20% | | 14-27% | 20% | 26% | |
| Social associations | 32.2 | | | 22.0 | 17.3 | |
| Violent crime | 48 |  | | 59 | 240 | |
| Injury deaths | 44 | | 26-70 | 50 | 64 | |
| Physical Environment | | | | | | 42 |
| Air pollution - particulate matter | 11.1 |  | | 9.5 | 10.0 | |
| Drinking water violations | 0% | | | 0% | 3% | |
| Severe housing problems | 7% | | 4-9% | 9% | 11% | |
| Driving alone to work | 78% | | 75-82% | 71% | 79% | |

| | Traill County | Trend(Click for info) | Error Margin | Top U.S. Performers* | | North Dakota | Rank (of 47) |
|--------------------------------------------------------------------------------------------------------|---------------|-----------------------|--------------|----------------------|-----|--------------|--------------|
| Long commute - driving alone | 29% | | 24-33% | 15% | 13% | | |
| * 90th percentile, i.e., only 10% are better. Note: Blank values reflect unreliable or missing data | | | | | | | 2015 |

2015

Steele County

| | Steele County | Trend(Click for info) | Error Margin | Top U.S. Performers* | North Dakota | Rank (of 47) |
|----------------------------------|---------------|-------------------------------------------------------------------------------------|--------------|----------------------|--------------|--------------|
| Health Outcomes | | | | | | NR |
| Length of Life | | | | | | NR |
| Premature death | | | | 5,200 | 6,388 | |
| Quality of Life | | | | | | NR |
| Poor or fair health | 12% | | 7-21% | 10% | 12% | |
| Poor physical health days | 1.9 | | 1.1-2.7 | 2.5 | 2.7 | |
| Poor mental health days | 2.3 | | 1.2-3.5 | 2.3 | 2.4 | |
| Low birthweight | | | | 5.9% | 6.5% | |
| Health Factors | | | | | | NR |
| Health Behaviors | | | | | | NR |
| Adult smoking | 12% | | 7-20% | 14% | 18% | |
| Adult obesity | 32% |  | 25-39% | 25% | 30% | |
| Food environment index | 7.2 | | | 8.4 | 8.5 | |
| Physical inactivity | 30% |  | 23-38% | 20% | 25% | |
| Access to exercise opportunities | 27% | | | 92% | 68% | |
| Excessive drinking | 26% | | 16-39% | 10% | 22% | |
| Alcohol-impaired driving deaths | 50% | | | 14% | 46% | |
| Sexually transmitted infections | | | | 138 | 416 | |
| Teen births | 21 | | | 20 | 28 | |
| Clinical Care | | | | | | NR |
| Uninsured | 12% |  | 11-14% | 11% | 12% | |
| Primary care physicians | 1,989:0 | | | 1,045:1 | 1,279:1 | |

| | Steele County | Trend(Click for info) | Error Margin | Top U.S. Performers* | North Dakota | Rank (of 47) |
|--------------------------------------|---------------|-----------------------|--------------|----------------------|--------------|--------------|
| Dentists | 1,960:0 | | | 1,377:1 | 1,710:1 | |
| Mental health providers | | | | 386:1 | 638:1 | |
| Preventable hospital stays | | | | 41 | 56 | |
| Diabetic monitoring | 87% | | 54-100% | 90% | 86% | |
| Mammography screening | 78.9% | | 39.0-100.0% | 70.7% | 68.0% | |
| Social & Economic Factors | | | | | | NR |
| High school graduation | | | | | 85% | |
| Some college | 72.3% | | 58.4-86.2% | 71.0% | 74.4% | |
| Unemployment | 2.2% | | | 4.0% | 2.9% | |
| Children in poverty | 12% | | 8-15% | 13% | 12% | |
| Income inequality | 3.8 | | 3.1-4.6 | 3.7 | 4.4 | |
| Children in single-parent households | 15% | | 6-25% | 20% | 26% | |
| Social associations | 10.1 | | | 22.0 | 17.3 | |
| Violent crime | 25 | | | 59 | 240 | |
| Injury deaths | | | | 50 | 64 | |
| Physical Environment | | | | | | NR |
| Air pollution - particulate matter | 10.7 | | | 9.5 | 10.0 | |
| Drinking water violations | 0% | | | 0% | 3% | |
| Severe housing problems | 6% | | 2-10% | 9% | 11% | |
| Driving alone to work | 75% | | 69-81% | 71% | 79% | |
| Long commute - driving alone | 24% | | 17-31% | 15% | 13% | |

* 90th percentile, i.e., only 10% are better. Note: Blank values reflect unreliable or missing data.



SANFORD[®]
HEALTH